

A Direct Approach for Sparse Quadratic Discriminant Analysis

Binyan Jiang, Xiangyu Wang, and Chenlei Leng *

September 21, 2016

Abstract

Quadratic discriminant analysis (QDA) is a standard tool for classification due to its simplicity and flexibility. Because the number of its parameters scales quadratically with the number of the variables, QDA is not practical, however, when the dimensionality is relatively large. To address this, we propose a novel procedure named QUDA for QDA in analyzing high-dimensional data. Formulated in a simple and coherent framework, QUDA aims to directly estimate the key quantities in the Bayes discriminant function including quadratic interactions and a linear index of the variables for classification. Under appropriate sparsity assumptions, we establish consistency results for estimating the interactions and the linear index, and further demonstrate that the misclassification rate of our procedure converges to the optimal Bayes risk, even when the dimensionality is exponentially high with respect to the sample size. An efficient algorithm based on the alternating direction method of multipliers (ADMM) is developed for finding interactions, which is much faster than its competitor in the literature. The promising performance of QUDA is illustrated via extensive simulation studies and the analysis of two datasets.

Key Words: *Bayes risk; Consistency; High-dimensional data; Linear discriminant analysis; Quadratic discriminant analysis; Sparsity.*

*Jiang is with Hong Kong Polytechnic University and Carnegie Mellon University. Wang is with Duke University. Leng is with University of Warwick. Corresponding author: Xiangyu Wang (xw56@stat.duke.edu).

1 Introduction

Classification is a central topic in statistical learning and data analysis. Due to its simplicity for producing quadratic decision boundaries, quadratic discriminant analysis (QDA) has become an important technique for classification, adding an extra layer of flexibility to the linear discriminant analysis (LDA); see Hastie, Tibshirani, and Friedman (2009). Despite its usefulness, the number of the parameters needed by QDA scales squarely with that of the variables, making it quickly inapplicable for problems with large or even moderate dimensionality. This problem is extremely eminent in the era of big data, as one often encounters datasets with the dimensionality larger, often times substantially larger than the sample size. This paper aims to develop a novel classification approach named QUDA to make QDA useful for analyzing ultra-high dimensional data.

For ease of presentation, we focus on binary problems where observations are from two classes. Suppose that the observations from class 1 follow $X \sim N(\mu_1, \Sigma_1)$ and those from class 2 satisfy $Y \sim N(\mu_2, \Sigma_2)$, where $\mu_k \in \mathbb{R}^p, k = 1, 2$ are the mean vectors and $\Sigma_k \in \mathbb{R}^{p \times p}, k = 1, 2$ are the two covariance matrices. Compared with LDA, it is assumed that $\Sigma_1 \neq \Sigma_2$ in QDA, which gives rise to a class boundary that is quadratic in terms of the variables. Bayes' rule classifies a new observation z to class 1 if $\pi_1 f(z|\mu_1, \Sigma_1) > \pi_2 f(z|\mu_2, \Sigma_2)$, where $f(z|\mu, \Sigma)$ is the probability density function of a multivariate normal distribution with mean μ and variance Σ , and π_1 and π_2 are the two prior probabilities. Following simple algebra, the Bayes discriminant function for a new observation z is seen as

$$D(z) = (z - \mu)^T \Omega (z - \mu) + \delta^T (z - \mu) + \eta,$$

where $\mu = (\mu_1 + \mu_2)/2$ is the mean of the two centroids, $\Omega = \Sigma_2^{-1} - \Sigma_1^{-1}$ is the difference of the two precision matrices, $\delta = (\Sigma_1^{-1} + \Sigma_2^{-1})(\mu_1 - \mu_2)$, and $\eta = 2 \log(\pi_1/\pi_2) + \frac{1}{4}(\mu_1 - \mu_2)^T \Omega (\mu_1 - \mu_2) + \log |\Sigma_2| - \log |\Sigma_1|$; see for example Anderson (2003). Note that in our setup, the discriminant function reduces to that of LDA when $\Sigma_1 = \Sigma_2 = \Sigma$. Completely

analogous to a two-way interaction model in linear regression, δ in the discriminant function $D(z)$ can be seen as a linear index of the variables whose nonzero entries play the role of main effects, whereas the nonzero entries in Ω can be understood as interactions of second-order between the variables. Note that there are other ways to represent the discriminant function. We choose to use $D(z)$ as it is a quadratic function of $z - \mu$, making the discriminant function location-invariant with respect to the coordinates. For easy reference, we shall call subsequently the parameters Ω, δ, μ , and η in the Bayes discriminant function collectively as Bayes components.

We highlight the main contributions of this paper as follows.

1. Our approach is the first *direct* approach for sparse QDA in a high-dimensional setup;
2. For estimating Ω , an intermediate step of QUDA and a problem of interest in its own right, we develop a new algorithm which is much more computationally and memory efficient than its competitor. See Section 2.1;
3. We develop new theory to show the theoretical attractiveness of the QUDA. In particular, the theory for estimating δ is new. See Section 3;
4. The problem of finding the right intercept η is of considerable interest. Since it is a scalar, a useful procedure is to choose η to minimize the empirical misclassification error once the other parameters are estimated. However, a general theory on estimated η as such is lacking (Hastie, Tibshirani, and Friedman, 2009). We provide a first theory for the convergence properties of our estimated η . See Section 3.4;
5. Empirically, the QUDA approach outperforms many of its competitors, especially when variables under consideration interact. See Section 4.

As more and more modern datasets are high-dimensional, the problem of classification in this context has received increasing attention as the usual practice of using empirical estimates for the Bayes components is no longer applicable. Bickel and Levina (2004) first highlighted that

LDA is equivalent to random guessing in the worst case scenario when the dimensionality is larger than the sample size. Scientifically and practically in many problems, however, the components in the Bayes discriminant function can be assumed sparse. In the problem we study in this paper, loosely speaking, the notion of sparsity entertains that the two-way interaction representation of the model only admits a small number of main effects and interactions. In the past few years, a plethora of methods built on suitable sparsity assumptions have been proposed to estimate the main effects as in LDA; see for example Shao et al. (2011), Cai and Liu (2011), Fan, Feng, and Tong (2012), Mai, Zou, and Yuan (2012), Mai and Zou (2013), and Jiang, Chen, and Leng (2015). Other related linear methods for high-dimensional classification can be found in Leng (2008), Witten and Tibshirani (2011), Pan, Wang and Li (2015), Mai, Yang, and Zou (2015), among others.

As pointed out by Fan et al. (2015b) and Sun and Zhao (2015), it has been increasingly recognized that the assumption of a common covariance matrix across different classes, needed by LDA, can be restrictive in many practical problems. The extra layer of flexibility offered by QDA that deals with two-way variable interactions makes it extremely attractive for such problems. Li and Shao (2015) studied sparse QDA by making sparsity assumptions on $\mu_2 - \mu_1$, Σ_1 , Σ_2 and $\Sigma_1 - \Sigma_2$ and proposed their sparse estimates. The assumptions made are not directly on the key quantities needed in the discriminant function $D(z)$. In addition, good estimates of these four quantities do not necessarily translate to better classification, a phenomenon similarly argued and observed by Cai and Liu (2011) and Mai, Zou, and Yuan (2012) for LDA. Fan et al. (2015b) proposed a screening method to identify interactions when Ω admits a two block sparse structure after permutation, before applying penalized logistic regression on the identified interactions and all the main effects to estimate a sparser model. Their method cannot deal with problems where the support of Ω is in general positions, for example, when Σ is a diagonal matrix. Further, the use of a separate second-step penalized logistic regression to determine important interactions and main effects is less appealing from a methodological perspective. Fan et al. (2015a) suggested a Rayleigh quotient based method for which all the fourth cross-moments of the predictors have to be estimated. Despite all

these efforts, a direct yet simple approach for QDA with less stringent assumptions than in Li and Shao (2015) for high-dimensional analysis is missing.

The proposed QUDA approach in this paper aims to overcome the difficulties mentioned above. In particular, compared with Li and Shao (2015), we only make sparsity assumptions on Ω and δ and estimate these two quantities directly in QUDA. Compared to Fan et al. (2015b), we allow the interactions in Ω in general positions, without resorting to a second stage approach for interactions and main effects selection. Compared with Fan et al. (2015a), we operate directly on QDA for which only second cross-moments of the variables are needed.

QUDA can also be understood as a novel attempt to select interactions in the discriminant function that correspond to the nonzero entries in Ω . The problem of interaction selection is a problem of its own importance and has been studied extensively recently for regression problems. See, for example, Bien, Taylor, and Tibshirani (2013), Hao and Zhang (2014), Hao, Feng, and Zhang (2014), Zhao and Leng (2015) and references therein. Despite the apparent similarity to linear regression with interactions, a key advantage of viewing QUDA as an interaction model is that $D(z)$ is automatically invariant with respect to centering and scaling of the variables, avoiding a long standing debate over importance of variables in regression (cf. Hao and Zhang (2015)). The problem of estimating Ω alone has also attracted attention lately in a different context. To understand how the structure of a network differs between different conditions and to find the common structures of two different Gaussian graphical models, Zhao, Cai, and Li (2014) proposed a direct approach for estimating Ω by formulating their procedure via the Dantzig selector. A severe limitation is that their linear programming procedure needs to deal with $O(p^2)$ constraints, and the memory requirement by the large constraint matrix is of the order $O(p^4)$. As a result, an iteration of the algorithm in Zhao, Cai, and Li (2014) requires $O(sp^4)$ computations, where s is the cardinality of the support of Ω . Apparently, their method does not scale well to high dimensional data. In Zhao, Cai, and Li (2014), problems with maximum size $p = 120$ were attempted and it was reported that a long time was needed to run their method. In contrast, we use a lasso formulation and develop a new algorithm based on the alternating direction methods of

multipliers (ADMM) for estimating Ω . The memory requirement of our algorithm is of the order $O(p^2)$ and its computational cost is of the order $O(p^3)$ per iteration, enabling QUDA to easily handle much larger problems.

The rest of the paper is organized as follows. Section 2 outlines the main QUDA methodology for estimating Ω and δ . A novel algorithm based on ADMM for estimating Ω is developed. Section 3 investigates the theory of QUDA and provides various consistency results for estimating Ω , δ , and η , as well as establishing the consistency of the misclassification risk relative to the Bayes risk. Section 4 presents extensive numerical studies and analysis of two datasets. Comparison with other classification methods demonstrates that QUDA is very competitive in estimating the sparsity pattern and the parameters of interest. We provide a short discussion and outline future directions of research in Section 5. All the Proof are relegated to the Supplementary Materials.

2 QUDA Methodology

To obtain an estimator for Bayes discriminant function $D(z)$, we propose direct estimators for the two of its Bayes components $\Omega = \Sigma_2^{-1} - \Sigma_1^{-1}$ and $\delta = (\Sigma_1^{-1} + \Sigma_2^{-1})(\mu_1 - \mu_2)$ under appropriate sparsity assumptions. Given data X_j , $j = 1, \dots, n_1$ from class 1 and Y_k , $k = 1, \dots, n_2$ from class 2, we can estimate μ_i and Σ_i , $i = 1, 2$, via their sample versions as

$$\hat{\mu}_1 = \frac{1}{n_1} \sum_{j=1}^{n_1} X_j, \quad \hat{\mu}_2 = \frac{1}{n_2} \sum_{j=1}^{n_2} Y_j;$$

$$\hat{\Sigma}_1 = \frac{1}{n_1} \sum_{j=1}^{n_1} (X_j - \hat{\mu}_1)(X_j - \hat{\mu}_1)^T, \quad \hat{\Sigma}_2 = \frac{1}{n_2} \sum_{j=1}^{n_2} (Y_j - \hat{\mu}_2)(Y_j - \hat{\mu}_2)^T.$$

When $p \gg \max\{n_1, n_2\}$, $\hat{\Sigma}_1$ and $\hat{\Sigma}_2$ are degenerate and cannot be directly used for estimating Ω . Denote the QUDA estimates of Ω as $\hat{\Omega}$ and δ as $\hat{\delta}$ which will be obtained as in (2) and (8) respectively. For a given scalar η , our QUDA procedure classifies z in to class 1

if

$$\left[z - \frac{\hat{\mu}_1 + \hat{\mu}_2}{2} \right]^T \hat{\Omega} \left[z - \frac{\hat{\mu}_1 + \hat{\mu}_2}{2} \right] + \hat{\delta}^T \left[z - \frac{\hat{\mu}_1 + \hat{\mu}_2}{2} \right] + \hat{\eta} > 0, \quad (1)$$

and classifies z into class 2 otherwise. From (1), we emphasize again that the nonzero entries in $\hat{\Omega}$ are the interactions of the variables that contribute to the classification rule, while the nonzero entries in $\hat{\delta}$ are the main effects of the variables that are used for classification. In the linear discriminant analysis when $\Sigma_1 = \Sigma_2$, the rule in (1) becomes the LDA rule which is linear in the variables. As $\eta = 2 \log(\pi_1/\pi_2) - \frac{1}{4}(\mu_1 - \mu_2)^T \Omega (\mu_1 - \mu_2) - \log |\Sigma_2| + \log |\Sigma_1|$ is a scalar, we can choose η as $\hat{\eta}$ using a simple grid search, bypassing the need especially to estimate the determinants of Σ_1 and Σ_2 . This is the strategy implemented in Section 4 and its analytical justification is provided in Section 3.4. Thus in the following, we shall focus on the estimation of Ω and δ , under certain sparsity assumptions on these two quantities.

2.1 Estimating Ω

Recall $\Omega = \Sigma_2^{-1} - \Sigma_1^{-1}$. It may be attempting to first estimate Σ_1^{-1} and Σ_2^{-1} as intermediate quantities before taking their difference. It is known, however, that accurate estimation of a covariance matrix or its inverse can be difficult in general in high dimensions unless additional assumptions are imposed (cf. Bickel and Levina (2008)). Because Ω is the quantity of interest that appears in the Bayes' rule, we propose to estimate it directly. To proceed, we note that $\Sigma_2 \Omega \Sigma_1 = \Sigma_1 \Omega \Sigma_2 = \Sigma_1 - \Sigma_2$. If we define a loss function as $Tr(\Omega^T \Sigma_1 \Omega \Sigma_2) / 2 - Tr(\Omega(\Sigma_1 - \Sigma_2))$, the loss function is minimized when Ω satisfies $\Sigma_2 \Omega \Sigma_1 = \Sigma_1 - \Sigma_2$ or $\Omega = \Sigma_2^{-1} - \Sigma_1^{-1}$. This simple observation motivates the following penalized loss formulation for estimating Ω by replacing $\Sigma_j, j = 1, 2$ by their empirical estimates as

$$\hat{\Omega} = \arg \min_{\Omega \in R^{p \times p}} \frac{1}{2} Tr(\Omega^T \hat{\Sigma}_1 \Omega \hat{\Sigma}_2) - Tr(\Omega(\hat{\Sigma}_1 - \hat{\Sigma}_2)) + \lambda \|\Omega\|_1, \quad (2)$$

where $\|\Omega\|_1$ is the ℓ_1 penalty of the vectorized Ω to encourage sparsity and λ is the tuning parameter. To obtain a symmetric estimator for Ω , we may simply use $\hat{\Omega}_0 = \frac{1}{2}(\hat{\Omega} + \hat{\Omega}^T)$ after $\hat{\Omega}$ is obtained. Because the second derivative of the above loss function is $\hat{\Sigma}_2 \otimes \hat{\Sigma}_1$ which is nonnegative definite, the formulation in (2) is a convex problem and can be solved by a convex optimization algorithm.

We now develop an ADMM algorithm to solve for $\hat{\Omega}$ in (2) (Boyd et al., 2011; Zhang and Zou, 2014). First write the optimization problem in (2) as

$$\min_{\Omega \in R^{p \times p}} \frac{1}{2} Tr \left(\Omega^T \hat{\Sigma}_1 \Omega \hat{\Sigma}_2 \right) - Tr \left(\Omega (\hat{\Sigma}_1 - \hat{\Sigma}_2) \right) + \lambda \|\Psi\|_1, \text{ s.t. } \Psi = \Omega. \quad (3)$$

From this, we can form the augmented Lagrangian as

$$L(\Omega, \Psi, \Lambda) = \frac{1}{2} Tr \left(\Omega^T \hat{\Sigma}_1 \Omega \hat{\Sigma}_2 \right) - Tr \left(\Omega (\hat{\Sigma}_1 - \hat{\Sigma}_2) \right) + \lambda \|\Psi\|_1 + Tr(\Lambda(\Omega - \Psi)) + \frac{\rho}{2} \|\Omega - \Psi\|_F^2,$$

where $\|\cdot\|_F$ is the Frobenius norm of a matrix and ρ is a parameter in the ADMM algorithm. See Section 4 for more details. Given the current estimate $\Omega^k, \Psi^k, \Lambda^k$, we update successively

$$\Omega^{k+1} = \arg \min_{\Omega \in R^{p \times p}} L(\Omega, \Psi^k, \Lambda^k), \quad (4)$$

$$\Psi^{k+1} = \arg \min_{\Psi \in R^{p \times p}} L(\Omega^{k+1}, \Psi, \Lambda^k), \quad (5)$$

$$\Lambda^{k+1} = \Lambda^k + \rho(\Omega^{k+1} - \Psi^{k+1}). \quad (6)$$

Write $\hat{\Sigma}_i = U_i D_i U_i^T$ as the eigenvalue decomposition of $\hat{\Sigma}_i$ where $D_i = \text{diag}(d_{i1}, \dots, d_{ip})$, $i = 1, 2$. Denote $A^k = (\hat{\Sigma}_1 - \hat{\Sigma}_2) - \Lambda^k + \rho \Psi^k$ and organize the diagonals of $(D_2 \otimes D_1 + \rho I)^{-1}$ in a matrix B , where $B_{jk} = 1/(d_{1j}d_{2k} + \rho)$. The following proposition provides explicit solutions for (4) and (5) which ensures efficient updation of our algorithm in each step.

Proposition 1. *Given Ψ^k, Λ^k, ρ and λ , the solution for (4) is given as:*

$$\Omega^{k+1} = U_1 [B \circ (U_1^T A^k U_2)] U_2^T;$$

Given Ω^{k+1}, Ψ^k and ρ , the solution for (5) is given as:

$$\Psi^{k+1} = S(\Omega^{k+1} + \frac{\Lambda^k}{\rho}, \frac{\lambda}{\rho}), \quad (7)$$

where S is known as the soft-thresholding operator on a matrix. Namely, the (i, j) entry of $S(A, b)$ for a matrix $A = (a_{ij})$ is $\text{sign}(a_{ij})(|a_{ij}| - b)_+$ where $(c)_+ = c$ for $c > 0$ and $(c)_+ = 0$ otherwise.

Note that for a given ρ , when updating Ω , we only need to update A^k which involves simple matrix subtraction, and then use matrix multiplication. The update in (4) can be efficiently implemented. Following is a brief derivation on how we obtain the expected solutions given in Proposition 1. For (4), note that the derivative of L with respect to Ω is

$$\hat{\Sigma}_1 \Omega \hat{\Sigma}_2 - (\hat{\Sigma}_1 - \hat{\Sigma}_2) + \Lambda^k + \rho(\Omega - \Psi^k) = (\hat{\Sigma}_1 \Omega \hat{\Sigma}_2 + \rho \Omega) - (\hat{\Sigma}_1 - \hat{\Sigma}_2) + \Lambda^k - \rho \Psi^k,$$

which can be written as

$$(\hat{\Sigma}_2 \otimes \hat{\Sigma}_1 + \rho I) \text{vec}(\Omega) = \text{vec} \left((\hat{\Sigma}_1 - \hat{\Sigma}_2) - \Lambda^k + \rho \Psi^k \right),$$

where vec is the vector operator. Using the equality $\text{vec}(AXB) = (B^T \otimes A) \text{vec}(X)$, we have

$$\text{vec}(\Omega) = (\hat{\Sigma}_2 \otimes \hat{\Sigma}_1 + \rho I)^{-1} \text{vec} \left((\hat{\Sigma}_1 - \hat{\Sigma}_2) - \Lambda^k + \rho \Psi^k \right).$$

and

$$(\hat{\Sigma}_2 \otimes \hat{\Sigma}_1 + \rho I)^{-1} = [(U_2 \otimes U_1)(D_2 \otimes D_1 + \rho I)(U_2^T \otimes U_1^T)]^{-1} = (U_2 \otimes U_1)(D_2 \otimes D_1 + \rho I)^{-1}(U_2^T \otimes U_1^T).$$

Then

$$\begin{aligned}
(\hat{\Sigma}_2 \otimes \hat{\Sigma}_1 + \rho I)^{-1} \text{vec}(A^k) &= [(U_2 \otimes U_1)(D_2 \otimes D_1 + \rho I)(U_2^T \otimes U_1^T)]^{-1} \text{vec}(A^k) \\
&= (U_2 \otimes U_1)(D_2 \otimes D_1 + \rho I)^{-1} (U_2^T \otimes U_1^T) \text{vec}(A^k) \\
&= (U_2 \otimes U_1)(D_2 \otimes D_1 + \rho I)^{-1} \text{vec}(U_1^T A^k U_2) \\
&= (U_2 \otimes U_1) \text{vec}(B \circ (U_1^T A^k U_2)) \\
&= \text{vec}(U_1 [B \circ (U_1^T A^k U_2)] U_2^T)
\end{aligned}$$

where $\circ$ is the Hadamard product. Therefore,

$$\Omega = U_1 [B \circ (U_1^T A^k U_2)] U_2^T.$$

Next we examine (5). Ignoring terms that are independent of Ψ , we just need to minimize

$$\frac{\rho}{2} \text{Tr}(\Psi^T \Psi) - \rho \text{Tr}((\Omega^{k+1})^T \Psi) - \text{Tr}((\Lambda^k)^T \Psi) + \lambda \|\Psi\|_1,$$

and the solution can be easily seen as (7). Again, the update for Γ can be efficiently implemented.

Our algorithm can be now summarized as following.

1. Initialize Ω , Ψ and Λ . Fix ρ . Compute the singular value decomposition $\hat{\Sigma}_1 = U_1 D_1 U_1^T$ and $\hat{\Sigma}_2 = U_2 D_2 U_2^T$, and compute B where $B_{jk} = 1/(d_{1j}d_{2k} + \rho)$. Repeat steps 2-4 until convergence;
2. Compute $A = (\hat{\Sigma}_1 - \hat{\Sigma}_2) - \Lambda + \rho \Psi$. Then update Ω as $\Omega = U_1 [B \circ (U_1^T A U_2)] U_2^T$;
3. Update Ψ as in (7) by soft-thresholding $\Omega + \frac{\Lambda}{\rho}$ elementwise by $\frac{\lambda}{\rho}$;
4. Update Λ by $\Lambda \leftarrow \Lambda + \rho(\Omega - \Psi)$ as in (6).

Note that the algorithm involves singular value decomposition of $\hat{\Sigma}_1$ and $\hat{\Sigma}_2$ only once. The

rest of the algorithm only involves matrix addition and multiplication. Thus, the algorithm is extremely efficient. Compared with Zhao, Cai, and Li (2014) whose algorithm has computational complexity of the order at least $O(p^4)$ and a memory requirement of $O(p^4)$, our algorithm has a memory requirement of the order $O(p^2)$ and computational complexity of $O(p^3)$. As a result, our method can handle much larger problems.

2.2 The linear index δ

After having estimated Ω as $\hat{\Omega}$, we discuss the estimation of the linear index $\delta = (\Sigma_1^{-1} + \Sigma_2^{-1})(\mu_1 - \mu_2)$. We develop a procedure that avoids estimating Σ_1^{-1} and Σ_2^{-1} . First note that

$$\Sigma_1\delta = \Sigma_1(\Sigma_1^{-1} + \Sigma_2^{-1})(\mu_1 - \mu_2) = \Sigma_1(2\Sigma_1^{-1} + \Omega)(\mu_1 - \mu_2) = 2(\mu_1 - \mu_2) + \Sigma_1\Omega(\mu_1 - \mu_2),$$

$$\Sigma_2\delta = \Sigma_2(\Sigma_1^{-1} + \Sigma_2^{-1})(\mu_1 - \mu_2) = \Sigma_2(2\Sigma_2^{-1} - \Omega)(\mu_1 - \mu_2) = 2(\mu_1 - \mu_2) - \Sigma_2\Omega(\mu_1 - \mu_2),$$

and

$$(\Sigma_1 + \Sigma_2)\delta = 4(\mu_1 - \mu_2) + (\Sigma_1 - \Sigma_2)\Omega(\mu_1 - \mu_2).$$

The last equation is the derivative of $\delta^T(\Sigma_1 + \Sigma_2)\delta/2 - \{4(\mu_1 - \mu_2) + (\Sigma_1 - \Sigma_2)\Omega(\mu_1 - \mu_2)\}^T\delta$. Motivated by this, we estimate δ by a direct method using the lasso regularization, similar to the one in Mai, Zou, and Yuan (2012), as

$$\hat{\delta} = \arg \min_{\delta \in R^p} \frac{1}{2}\delta^T(\hat{\Sigma}_1 + \hat{\Sigma}_2)\delta - \hat{\gamma}^T\delta + \lambda_\delta\|\delta\|_1, \quad (8)$$

where $\hat{\gamma} = 4(\hat{\mu}_1 - \hat{\mu}_2) + (\hat{\Sigma}_1 - \hat{\Sigma}_2)\hat{\Omega}(\hat{\mu}_1 - \hat{\mu}_2)$, $\|\cdot\|_1$ is the vector ℓ_1 penalty and λ_δ is a tuning parameter. The optimization in (8) is a standard lasso problem and is easy to solve using existing lasso algorithms. We remark that (8) is much more challenging to analyze theoretically than the method in Mai, Zou, and Yuan (2012), since the accuracy of $\hat{\Omega}$ as an estimator of Ω has to be carefully quantified in $\hat{\gamma}$.

We emphasize that our framework is extremely flexible and can accommodate additional

constraints. As a concrete example, let's consider enforcing the so-called strong heredity principle in that an interaction is present unless the corresponding main effects are both present, i.e. if $\Omega_{jk} \neq 0$ then $\delta_j \neq 0$ and $\delta_k \neq 0$; see for example Hao and Zhang (2015). Denote $\mathcal{I} \subset \{1, \dots, p\}$ as the set such that for any $j, k \in \mathcal{I}$ there exists some $\hat{\Omega}_{jk} \neq 0$. We can change the penalty in (8) as $\|\delta_{\mathcal{I}^c}\|_1$ such that the variables in $\mathcal{I}$ are not penalized. Due to space limitation, this line of research will not be studied in the current paper.

3 Theory

We show that our method can consistently select the true nonzero interaction terms in Ω and the true nonzero terms in δ . In addition, we provide explicit upper bounds for the estimation error under l_∞ norm. For classification, we further show that the misclassification rate of our QUDA rule converges to the optimal Bayes risk under some sparsity assumptions. For simplicity in this section we assume that $n_1 \asymp n_2$ and write $n = \min\{n_1, n_2\} - 1$. Instead of assuming $\mu_2 - \mu_1$, Σ_1, Σ_2 and $\Sigma_1 - \Sigma_2$ to be sparse as in Li and Shao (2015), we only assume that Ω and δ are sparse. For the estimation of Ω , the rate in Corollary 1 is similar to the one in Theorem 3 of Zhao, Cai, and Li (2014). However, as we pointed out previously, our method is computationally much more efficient and scales better to large-dimensional problems. In addition, our work is the first direct estimation approach for sparse QDA. Importantly, the results for estimating δ are new.

Note that when estimating δ as in (8), we have used $\hat{\Omega}$ as a plug-in estimator for Ω . Consequently, from Corollaries 1 and 2, the error rate of $\hat{\delta}$ in estimating δ is a factor times of that of $\hat{\Omega}$ in estimating Ω . However, in the QUDA discriminant function defined as in (1), $\hat{\Omega}$ appears in the first term which is a product of three components while $\hat{\delta}$ appears in the second term which is a product of two components. As a consequence, the overall estimation error rates of these two terms become equal. This implies that even though the estimating error of $\hat{\Omega}$ might aggregate in the estimation of δ , it does not affect the convergence rate of the classification error at all. Below we provide theory for estimating Ω , δ , and η , as well as

quantifying the overall misclassification error rate.

3.1 Theory for estimating Ω

We first introduce some notation. We assume that $\Omega = (\Omega_{ij})_{1 \leq i, j \leq p}$ is sparse with support $S = \{(i, j) : \Omega_{ij} \neq 0\}$ and we use S^c to denote the complement of S . Let d be the maximum node degree in Ω . For a vector $x = (x_1, \dots, x_p)^T$, the l_q norm is defined as $|x|_q = (\sum_{i=1}^p |x_i|^q)^{1/q}$ for any $1 \leq q < \infty$ and the l_∞ norm is defined as $|x|_\infty = \max_{1 \leq i \leq p} |x_i|$. For any matrix $M = (m_{ij})_{1 \leq i, j \leq p}$, its entrywise l_1 norm is defined as $\|M\|_1 = \sum_{1 \leq i, j \leq p} |m_{ij}|$ and its entrywise l_∞ norm is written as $\|M\|_\infty = \max_{1 \leq i, j \leq p} |m_{ij}|$. We use $\|M\|_{1,\infty} = \max_{1 \leq i \leq p} \sum_{j=1}^p |m_{ij}|$ to denote the l_1/l_∞ norm induced matrix-operator norm. We denote $\Gamma = \Sigma_2 \otimes \Sigma_1$ and $\hat{\Gamma} = \hat{\Sigma}_2 \otimes \hat{\Sigma}_1$. Write $\Sigma_k = (\sigma_{kij})_{1 \leq i, j \leq p}$, $\hat{\Sigma}_k = (\hat{\sigma}_{kij})_{1 \leq i, j \leq p}$ for $k = 1, 2$. By the definition of Kronecker product, Γ is a $p^2 \times p^2$ matrix indexed by vertex pairs in that $\Gamma_{(i,j),(k,l)} = \sigma_{1ik}\sigma_{2jl}$. Denote $\Delta_i = \hat{\Sigma}_i - \Sigma_i$ for $i = 1, 2$, $\Delta_\Gamma = \hat{\Gamma} - \Gamma$, $\Delta_{\Gamma^T} = \hat{\Gamma}^T - \Gamma^T$, and $\epsilon_i = \|\Delta_i\|_\infty$, $\epsilon = \max\{\epsilon_1, \epsilon_2\}$. $B = \max\{\|\Sigma_1\|_\infty, \|\Sigma_2\|_\infty\}$, $B_\Sigma = \max\{\|\Sigma_1\|_{1,\infty}, \|\Sigma_2\|_{1,\infty}\}$ and $B_\Gamma = \|\Gamma_{S,S}^{-1}\|_{1,\infty}$, $B_{\Gamma^T} = \|(\Gamma_{S,S}^T)^{-1}\|_{1,\infty}$, $B_{\Gamma,\Gamma^T} = \max\{B_\Gamma, B_{\Gamma^T}\}$.

To establish the model selection consistency of our estimator, we assume the following irrepresentability condition:

$$\alpha = 1 - \max_{e \in S^c} |\Gamma_{e,S} \Gamma_{S,S}^{-1}|_1 > 0.$$

This condition was first introduced by Zhao and Yu (2006) to establish the model selection consistency of the lasso. The following theorem gives the model selection consistency and the rate of convergence for the estimation of Ω .

Theorem 1. *Assume that $\alpha > 0$. By choosing*

$$\lambda = \max \left\{ 8\alpha^{-1}, \frac{3(2-\alpha)(2B+1)}{1-\alpha} d^2 B B_{\Gamma,\Gamma^T} [1 + 2(B_\Sigma^2 + 1/3) B_{\Gamma,\Gamma^T}] \right\} \sqrt{\frac{c \log p + \log C_1}{C_2 n}},$$

for some $c > 2$ and

$$n > \max\{(C_2 \min(B^2, 1))^{-1}, 81B^2d^4B_{\Gamma, \Gamma^T}^2C_2^{-1}, 9(C_2\alpha)^{-1}B^2B_{\Gamma, \Gamma^T}^2[1 + 2(B_\Sigma^2 + 1/3)B_{\Gamma, \Gamma^T}]^2\} \\ \cdot (c \log p + \log C_1),$$

we have, with probability greater than $1 - p^{2-c}$,

$$(i) \hat{\Omega}_{S^c} = 0;$$

$$(ii)$$

$$\|\hat{\Omega} - \Omega\|_\infty < C_3 \sqrt{\frac{c \log p + \log C_1}{C_2 n}},$$

$$\text{where } C_3 = \frac{14}{3} \max \left\{ 8\alpha^{-1}, \frac{3(2-\alpha)(2B+1)}{1-\alpha} d^2 B B_{\Gamma, \Gamma^T} [1 + 2(B_\Sigma^2 + 1/3) B_{\Gamma, \Gamma^T}] \right\} B_{\Gamma, \Gamma^T} + 24d^2 B^2 B_{\Gamma, \Gamma^T}^2.$$

Here C_1 and C_2 are constants defined as in Lemma 1. Theorem 1 states that if the irrepresentability condition is satisfied, the support of Ω is estimated consistently, and the rate of convergence of estimating Ω under l_∞ norm is of order $O\left(C_3 \sqrt{\frac{\log p}{n}}\right)$, where C_3 depends on the sparsity Σ_1, Σ_2 and their Kronecker product. In particular, from the proof of Theorem 1, we have the following corollary.

Corollary 1. Assume that $\alpha > 0$, $B_\Sigma < \infty$ and $B_{\Gamma, \Gamma^T} < \infty$. For any constant $c > 2$, choosing $\lambda = Cd^2 \sqrt{\frac{\log p}{n}}$ for some constant $C > 0$, if $d^2 \sqrt{\frac{\log p}{n}} \rightarrow 0$, we have with probability greater than $1 - p^{2-c}$, $\hat{\Omega}_{S^c} = 0$ and

$$\|\hat{\Omega} - \Omega\|_\infty = O\left(d^2 \sqrt{\frac{\log p}{n}}\right).$$

Similar to Condition 2 in Zhao, Cai, and Li (2014), the assumption $B_{\Gamma, \Gamma^T} < \infty$ in Corollary 1 is closely related to the mutual incoherence property introduced in Donoho and Huo (2001). In fact, it holds when imposing the usual mutual incoherence condition on the inverses of the submatrices (indexed by S) of Σ_1 and Σ_2 . Since d is the maximum node

degree in Ω , the number of nonzero entries in Ω is of order $O(dp)$. the rate $O\left(d^2\sqrt{\frac{\log p}{n}}\right)$ we obtained in Corollary 1 is better than the rate in Theorem 3 of Zhao, Cai, and Li (2014). However, in the case where only $O(d)$ covariates and some of their interactions are important, our rate is the same as the one in Zhao, Cai, and Li (2014).

3.2 Theory for estimating δ

Let $D = \{i : \delta_i \neq 0\}$ be the support of δ and let d_δ be its cardinality. Denote $A_1 = \|\Omega\|_{1,\infty}$, $A_2 = \|(\Omega^{-1})_{\cdot,D}\|_{1,\infty}$, $\epsilon_\mu = \max\{|\mu_1 - \hat{\mu}_1|_\infty, |\mu_2 - \hat{\mu}_2|_\infty\}$. We define $A_\Sigma = \|\Sigma_{D,D}^{-1}\|_{1,\infty}$ and write $\hat{\Sigma} = (\hat{\Sigma}_1 + \hat{\Sigma}_2)/2$, $\gamma = 4(\mu_1 - \mu_2) + (\Sigma_1 - \Sigma_2)\Omega(\mu_1 - \mu_2)$, $\Delta_\mu = \mu_1 - \mu_2$, $\hat{\Delta}_\mu = \hat{\mu}_1 - \hat{\mu}_2$, $A_\gamma = \|\gamma\|_\infty$ and $\|\hat{\Omega} - \Omega\|_\infty = \epsilon_\Omega$.

Let $\Sigma = (\Sigma_1 + \Sigma_2)/2$. To establish the model selection consistency of our estimator $\hat{\delta}$, we assume the following irrepresentability condition:

$$\alpha_\delta = 1 - \max_{e \in S^c} |\Sigma_{e,D} \Sigma_{D,D}^{-1}|_1 > 0.$$

The following theorem gives the model selection consistency and the rate of convergence for the estimation of δ .

Theorem 2. *Under the same assumptions in Theorem 1 and assuming that*

$$\begin{aligned} \lambda_\delta = & \max \left\{ \frac{2(2 - \alpha_\delta)}{\alpha_\delta} [4 + (2 + B_\Sigma A_2) |\Omega(\mu_1 - \mu_2)|_1 + 2B(A_1 + C_3)], \frac{d_\delta(5A_\Sigma + 2B_\Sigma A_\Sigma^2)}{C_\delta \alpha_\delta} \right\} \\ & \cdot (C_3 + 1) \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}}, \end{aligned}$$

and

$$\begin{aligned} n > & \max \{ C_{2\delta}^{-1}, 2C_2^{-1} A_\Sigma^2 d_\delta^2, C_2^{-1} (A_2 + 1)^2 d^2, C_2^{-1} C_\delta^{-2} \alpha_\delta^{-2} d_\delta^2 (5A_\Sigma + 2B_\Sigma A_\Sigma^2)^2 \} \\ & \cdot (c \log p + \log C_1), \end{aligned}$$

where $0 < C_\delta = \frac{\alpha_\delta \lambda_\delta - (2 - \alpha_\delta) K_\gamma}{\alpha_\delta \lambda_\delta (1 + A_\gamma + K_\gamma)} < 1$ and $C_{2\delta} = \min\{C_2, (2\sigma^2)^{-1}\} \cdot \min\{B^2, 1\}$, we have with probability greater than $1 - p^{2-c}$,

$$(i) \hat{\delta}_{D^c} = 0;$$

$$(ii)$$

$$\|\hat{\delta} - \delta\|_\infty < C_{3\delta} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}},$$

where

$$\begin{aligned} C_{3\delta} = & 2A_\gamma A_\Sigma^2 d_\delta + 4A_\Sigma [4 + (2 + B_\Sigma A_2) |\Omega(\mu_1 - \mu_2)|_1 + 2B(A_1 + C_3)] (C_3 + 1) + 2A_\Sigma (C_3 + 1) \\ & \cdot \max \left\{ \frac{2(2 - \alpha_\delta)}{\alpha_\delta} [4 + (2 + B_\Sigma A_2) |\Omega(\mu_1 - \mu_2)|_1 + 2B(A_1 + C_3)], \frac{d_\delta (5A_\Sigma + 2B_\Sigma A_\Sigma^2)}{C_\delta \alpha_\delta} \right\}. \end{aligned}$$

From Theorem 2 and Corollary 1 we immediately have:

Corollary 2. Let $d_0 = \max\{d, d_\delta\}$. Under the assumptions of Corollary 1 and assume that $\alpha_\delta > 0$, $A_\gamma < \infty$, $A_\Sigma < \infty$ and $A_1, A_2 < \infty$, $|\Omega(\mu_1 - \mu_2)|_\infty = O(d_0^2)$. For any constant $c > 0$, by choosing $\lambda = C d_0^2 \sqrt{\frac{\log p}{n}}$ for some constant $C > 0$, and assume that $d_0^3 \sqrt{\frac{\log p}{n}} \rightarrow 0$, we have with probability greater than $1 - p^{2-c}$,

$$\|\hat{\delta} - \delta\|_\infty = O \left(d_0^3 \sqrt{\frac{\log p}{n}} \right).$$

When $\Sigma_1 = \Sigma_2$, δ reduces to $2\Sigma_1^{-1}(\mu_1 - \mu_2)$, which is proportional to the direct discriminant variable β in Mai, Zou, and Yuan (2012), Cai and Liu (2011) and Fan, Feng, and Tong (2012), and variables in $D = \{i : \delta_i \neq 0\}$ are linear discriminative features contributing to the Bayes rule. From the proof of Theorem 2 and the rate given in Theorem 2, we can see that when $A_\gamma < \infty$, $\|\hat{\delta} - \delta\|_\infty$ is of order $O(A_\Sigma \lambda_\delta)$. This is consistent to the result obtained in Theorem 1 of Mai, Zou, and Yuan (2012) for the $\Sigma_1 = \Sigma_2$ case.

3.3 Misclassification rate

In this subsection, we study the asymptotic behavior of the misclassification rate for a given η and postpone the theory when η is estimated to Section 3.4. Let $R(i|j)$ and $R_n(i|j)$ be the probabilities that a new observation from class j is misclassified to class i by Bayes' rule and the QUDA rule respectively. Suppose $2 \log(\pi_1/\pi_2) = \eta - \frac{1}{4}(\mu_1 - \mu_2)^T \Omega(\mu_1 - \mu_2) - \log |\Sigma_2| + \log |\Sigma_1|$. The optimal Bayes risk is given as

$$R = \pi_1 R(2|1) + \pi_2 R(1|2),$$

and the misclassification rate of the QUDA rule takes the following form:

$$R_n = \pi_1 R_n(2|1) + \pi_2 R_n(1|2).$$

Suppose $z_i \sim N(\mu_i, \Sigma_i)$ for $i = 1, 2$. Denote the density of $(z_i - \mu)^T \Omega(z_i - \mu) + \delta^T(z_i - \mu) + \eta$ as $F_i(z)$. For any constant c , define

$$u_c = \max\{\text{ess sup}_{z \in [-c, c]} F_i(z), i = 1, 2\},$$

where ess sup denotes the essential supremum which is defined as supremum on almost everywhere of the support, i.e., except on a set of measure zero. Let s be the number of nonzero elements in Ω . The following theorem establishes upper bounds for the misclassification rate difference between R_n and R .

Theorem 3. *Assuming that there exist constants $C_\mu > 0, C_\Sigma > 1$ such that $\max\{|\mu_1|_\infty, |\mu_2|_\infty\} \leq C_\mu$, and $C_\Sigma^{-1} \leq \min\{\lambda_p(\Sigma_1), \lambda_p(\Sigma_2)\} \leq \max\{\lambda_1(\Sigma_1), \lambda_1(\Sigma_2)\} \leq C_\Sigma$ where $\lambda_i(\Sigma_j)$ denotes the i th largest eigenvalue of Σ_j . Under the assumptions of Theorems 1 and 2, we have:*

(i) if $sC_3 \sqrt{\frac{c \log p + \log C_1}{C_2 n}} + d_\delta C_{3\delta} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}} \rightarrow 0$ and there exist positive constants c, U_c

such that $u_c \leq U_c < \infty$, then

$$R_n - R = O_p \left(sC_3 \sqrt{\frac{c \log p + \log C_1}{C_2 n}} + d_\delta C_{3\delta} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}} \right);$$

(ii) if $\left(sC_3 \log p \sqrt{\frac{c \log p + \log C_1}{C_2 n}} + d_\delta C_{3\delta} \sqrt{\log p} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}} \right) (1 + u_c) \rightarrow 0$ for some positive constant c , then with probability greater than $1 - 3p^{2-c}$ for some constant $c > 2$,

$$R_n - R = O \left(\left(sC_3 \log p \sqrt{\frac{c \log p + \log C_1}{C_2 n}} + d_\delta C_{3\delta} \sqrt{\log p} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}} \right) (1 + u_c) \right).$$

Theorem 3 (i) indicates that under appropriate sparsity assumptions, our QUDA rule is optimal in that its misclassification rate converges to the optimal Bayes risk in probability. The second statement of Theorem 3 states that under stronger conditions, R_n converges to R with overwhelming probability. From Corollary 1 and Corollary 2 and the above theorem, we immediately have:

Corollary 3. *Under the assumptions of Corollary 1, Corollary 2 and Theorem 3, we have,*

(i) if $sd_0^2 \sqrt{\frac{\log p}{n}} \rightarrow 0$ and there exist positive constants c, U_c such that $u_c \leq U_c < \infty$, then

$$R_n - R = O_p \left(sd_0^2 \sqrt{\frac{\log p}{n}} \right);$$

(ii) if $(1 + u_c)sd_0^2 \sqrt{\frac{\log^3 p}{n}} \rightarrow 0$ for some constant $c > 0$, then with probability greater than $1 - 3p^{2-c}$ for some constant $c > 2$,

$$R_n - R = O \left(sd_0^2 (1 + u_c) \sqrt{\frac{\log^3 p}{n}} \right).$$

We remark that our assumption that $u_c \leq U_c$ for some constants c and U_c is similar to Condition (C4) in Li and Shao (2015), and our assumption is weaker in that we only assume

the densities $F_i(z)$ is bounded in a neighborhood of zero while Condition (C4) in Li and Shao (2015) states that the densities are bounded everywhere.

3.4 Choice of η

The question of choosing the scalar η is critical for classification but receives little attention in existing literature; see Mai, Zou, and Yuan (2012) for a detailed discussion for the LDA case. In this section, we propose to choose η by minimizing the in-sample misclassification error and establish analytical results for the estimation of η and the misclassification rate. With some abuse of notation, let (z_i, l_i) be our data where z_i 's are the covariates and l_i 's are the labels, i.e., $l_i \in \{0, 1\}$. To obtain $\hat{\eta}$, we seek to minimize the in-sample misclassification error given $\hat{\mu}$, $\hat{\delta}$ and $\hat{\Omega}$:

$$\hat{\eta} = \arg \min_e \frac{1}{n} \sum_{i=1}^n |I\{\hat{D}(z_i, e) > 0\} - l_i|,$$

where $\hat{D}(z, e) = (z - \hat{\mu})^T \hat{\Omega}(z - \hat{\mu}) + \hat{\delta}^T(z - \hat{\mu}) + e$. We write $d(z) = (z - \mu)^T \Omega(z - \mu) + \delta^T(z - \mu)$ and $\hat{d}(z) = (z - \hat{\mu})^T \hat{\Omega}(z - \hat{\mu}) + \hat{\delta}^T(z - \hat{\mu})$ and hence we have

$$\hat{D}(z_i, e) = \hat{d}(z_i) + e.$$

Then the object function becomes

$$\frac{1}{n} \sum_{i=1}^n |I\{\hat{d}(z_i) + e > 0\} - l_i|.$$

Without loss of generality, we can assume $\hat{d}(z_1) < \hat{d}(z_2) < \dots < \hat{d}(z_n)$. For any e we define the index $k(e)$ to be the largest $\hat{d}(z_k)$ that satisfies $\hat{d}(z_k) < -e < \hat{d}(z_{k+1})$. Thus, the

optimization can be further simplified as

$$\hat{\eta} = \arg \min_e \frac{1}{n} \left[\sum_{i=1}^{k(e)} l_i + \sum_{i=k(e)+1}^n (1 - l_i) \right].$$

Solving the above problem is simple. One just needs to compute the values of the object function for $k = 0, 1, 2, \dots, n$ and find the index k^* that minimizes its value. The optimal $\hat{\eta}$ can then be found as any value satisfying

$$\hat{\eta} \in (-\hat{d}(z_{k^*+1}), -\hat{d}(z_{k^*})).$$

Next we establish the asymptotic results for $\hat{\eta}$ and the misclassification rate. For a given e , we use $R(d, e)$ and $R(\hat{d}, e)$ to denote the misclassification rate associated with discriminant function $D(z, e) = d(z) + e$ and discriminant function $\hat{D}(z, e) = \hat{d}(z) + e$ respectively. Analogously, the in-sample misclassification rate of $D(z, e)$ and $\hat{D}(z, e)$ are denoted as $R_n(d, e)$ and $R_n(\hat{d}, e)$. From the optimality of the Bayes rule, we know that $\eta = 2 \log(\pi_1/\pi_2) + \frac{1}{4}(\mu_1 - \mu_2)^T \Omega(\mu_1 - \mu_2) + \log |\Sigma_2| - \log |\Sigma_1|$ is the unique minimizer of $R(d, \cdot)$ and we denote the corresponding optimal Bayes misclassification rate as $R = R(d, \eta)$. On the other hand, $\hat{\eta}$ is a minimizer of $R_n(\hat{d}, e)$. In order to make the estimation problem feasible, we assume that there exists a constant c such that $|\eta| < c < \infty$. The following proposition indicates that, although the 0-1 loss used for computing the misclassification rate is neither continuous nor convex, the misclassification rate has a desirable property.

Proposition 2. *$R(d, e)$ is strictly monotone increasing in $e \in [\eta, \infty)$ and strictly monotone decreasing in $e \in (-\infty, \eta]$.*

From Proposition 2 and following Theorem 5.7 of Van der Vaart (2011), we establish the following theorem, which indicates that the estimator $\hat{\eta}$ is consistent and the resulting misclassification rate using the estimated rule $\hat{D}(z, \hat{\eta})$ tends to the optimal Bayes misclassification rate in probability.

Theorem 4. *Let $\hat{\eta}$ be a minimizer of $R_n(\hat{d}, e)$. Under the assumptions of Theorem 3, we have*

(i) $\hat{\eta} \rightarrow \eta$ in probability;

(ii) $R_n(\hat{d}, \hat{\eta}) \rightarrow R$ in probability.

4 Numerical Study

In this section, we provide extensive numerical evidence to show the empirical performance of QUDA by comparing it to its competitors, including the sparse QDA (sQDA, Li and Shao (2015)), the innovated interaction screening for sparse quadratic discriminant analysis (IIS-SQDA, Fan et al. (2015b)), penalized logistic regression with only main effects considered (PLR), penalized logistic regression with all interaction terms (PLR2), the direct approach for sparse LDA (DSDA, Mai, Zou, and Yuan (2012)), the conventional LDA (LDA), the conventional QDA (QDA) and the oracle procedure (Oracle). The oracle procedure uses the true underlying model and serves as the optimal risk bound for comparison. We evaluate all methods via five synthetic datasets and two real datasets.

To fit QUDA, we employ ADMM to estimate Ω and the coordinate-wise descent algorithm (Friedman et al., 2010) to fit δ . Once Ω and δ are given, we then find the value of η by a simple linear search, minimizing the in-sample misclassification error. The rate parameter ρ in ADMM is set according to the optimal criterion suggested by Ghadimi et al. (2015). The other two tuning parameters, λ for estimating Ω and λ_δ for estimating δ , are chosen by 5-fold cross-validation, where the loss function is chosen to be the out-of-sample misclassification rate. We implemented sQDA in Matlab with the leave-one-out-cross-validation (Li and Shao, 2015) to tune the three parameters. We employ Matlab’s built-in function *fitcdiscr* to fit LDA and QDA and the R package *dsda* (Mai, Zou, and Yuan, 2012) to fit DSDA. For PLR, PLR2 and the second stage fit of IIS-SQDA which is a penalized logistic regression, we use the *glmnet* package and set $\alpha = 0.5$ as the elastic net parameter. Other values of α was

tried but did not change the result much. The other tuning parameter in *glmnet* is chosen by 10-fold cross-validation to minimize out-of-sample classification error. For the first stage of *IIS-SQDA* which is a screening step, we adopt the oracle-assisted approach proposed in Fan et al. (2015b), i.e., using the true Σ_1 and Σ_2 to compute the transformed variables used for screening as discussed in Fan et al. (2015b). To seek an appropriate screening size, we preserve the top 10, 30 or 50 variables for each experiment to form interaction terms and report the best result (smallest misclassification error) for *IIS-SQDA*.

4.1 Synthetic data

For synthetic data, we use the same setup in Fan et al. (2015b). Observations are simulated from $N(u_1, \Sigma_1)$ and $N(u_2, \Sigma_2)$ where $u_2 = 0$. Recall $\Omega_1 = \Sigma_1^{-1}$ and $\Omega_2 = \Sigma_2^{-1}$. We set $u_1 = \Sigma_1 \beta$ for $\beta = (0.6, 0.8, 0, \dots, 0)^T$. We consider three different dimensions $p = 50, 200$, or 500 with $n_1 = n_2 = 100$. The parameters Ω_1, Ω_2 and β are set as follows.

- Model 1: This model is Model 3 in Fan et al. (2015b) where Ω_1 is a band matrix with $(\Omega_1)_{ii} = 1$ and $\Omega_{ij} = 0.3$ for $|i - j| = 1$. We set $\Omega_2 = \Omega_1 + \Omega$, where Ω is a symmetric and sparse matrix with $\Omega_{10, 10} = -0.3758, \Omega_{10, 30} = 0.0616, \Omega_{10, 50} = 0.2037, \Omega_{30, 30} = -0.5482, \Omega_{30, 50} = 0.0286$, and $\Omega_{50, 50} = -0.4614$. The other 3 nonzero entries in the lower triangle of Ω are determined by symmetry.
- Model 2: We set $(\Omega_1)_{ij} = 0.5^{|i-j|}$ and let $\Omega_2 = \Omega_1 + \Omega$, where $\Omega = I_p$.
- Model 3: Ω_1 is the same as Model 2 and $\Omega_2 = \Omega_1$.
- Model 4: Ω_1 is the same as Model 2 and Ω is a band matrix defined as $(\Omega)_{ii} = 1$ and $(\Omega)_{ij} = 0.5$ for $|i - j| = 1$. Let $\Omega_2 = \Omega_1 + \Omega$.
- Model 5: $\Omega_1 = I_p$ and $\Omega_2 = \Omega_1 + \Omega$ where Ω is a random sparse symmetric matrix with conditional number 10 and non-zero density $n_1/p^2 \times 0.7$ (generated by *sprandsym* in Matlab).

Model 1 is a model where Ω is a sparse two-block matrix after permutation. This is a model that favors IIS-SQDA. In Model 2, the difference between Ω_1 and Ω_2 is a diagonal matrix, and IIS-SQDA is expected to underperform as its screening step for identifying variables that are involved in interaction would retain all the variables. Model 3 is obviously a model that favors the linear discriminant analysis (LDA) as $\Omega = 0$, and in particular favors the sparse LDA (DSDA). This model is simulated to test whether methods designed for sparse QDA work satisfactorily in situations where LDA works the best. In Model 4, the difference matrix Ω is a tridiagonal matrix where the screening step of IIS-SQDA is expected to underperform. Finally, in Model 5, Ω admits a random sparse structure having $0.7n_1 = 70$ nonzero entries regardless of the dimension p . Our implementation of IIS-SQDA is applied only to Model 2, 3, 4 and 5 while the results for Model 1 are directly cited from Fan et al. (2015b).

We simulate 100 synthetic datasets for each model and record for each method under comparison: 1) The misclassification rate (MR), 2) The false positives for main effects and interactions (FP.main for δ and FP.inter for Ω), 3) The false negatives for main effects and interactions (FN.main for δ and FN.inter for Ω). The results are summarized in Table 1, 2, 3, 4 and 5 for the five models under consideration. From these tables, we can observe the following phenomena.

1. For Model 1 where the setup favors IIS-SQDA, IIS-SQDA performs the best in terms of variable selection. QUDA performs similarly in terms of the misclassification rate. These two methods outperform LDA and QDA (when $p = 50$), and PLR, PLR2, DSDA by a large margin in classification error.
2. For Model 2, as expected, IIS-SQDA is outperformed by QUDA by a large margin. Interestingly, PLR2 performs the second best. Linear classifiers including PLR and DSDA perform quite badly.
3. For Model 3, DSDA, IIS-QDA and QUDA perform best and similarly. It is interesting that QUDA performs on par with DSDA even when the model clearly favors sparse linear classifiers.

4. For Model 4, QUDA outperforms all other methods again by a large margin.
5. For Model 5, QUDA performs the best and by a large margin when p becomes large.

To summarize, QUDA achieves the smallest misclassification rate in most examples and competitive performance in selecting main and interaction effects. IIS-SQDA is the preferred approach if Ω is a two-block diagonal matrix after permutation as is the case for Model 1. PLR2 generally performs better than (sparse) linear classifiers when interactions exist.

Table 1: The means and standard errors (in parentheses) of various performance measures by different classification methods for model 1 based on 100 replications

p	Method	MR (%)	FP.main	FP.inter	FN.main	FN.inter
50	LDA	39.43 (0.15)	—	—	—	—
	QDA	43.47 (0.10)	—	—	—	—
	PLR	36.12 (0.26)	5.95 (0.93)	—	1.21 (0.04)	—
	DSDA	35.05 (0.22)	8.81 (1.06)	—	0.07 (0.03)	—
	sQDA	27.64 (0.22)	11.17 (1.49)	—	0.33 (0.05)	—
	PLR2	30.15 (0.44)	0.51 (0.14)	11.26 (2.78)	0.60 (0.05)	2.62 (0.09)
	IIS-SQDA	27.56 (0.27)	5.60 (0.82)	2.16 (0.32)	0.19 (0.04)	2.05 (0.09)
	QUDA	26.50 (0.28)	0.85 (0.18)	35.26 (4.72)	0.39 (0.07)	3.74 (0.14)
	Oracle	23.04 (0.09)	—	—	—	—
200	PLR	37.62 (0.34)	7.82 (1.87)	—	1.47 (0.05)	—
	DSDA	36.34 (0.30)	15.06 (3.37)	—	0.36 (0.05)	—
	sQDA	26.80 (0.21)	12.75 (2.22)	—	0.47 (0.05)	—
	PLR2	32.55 (0.53)	0.25 (0.06)	17.44 (3.63)	0.90 (0.05)	2.72 (0.08)
	IIS-SQDA	26.94 (0.31)	6.43 (1.24)	0.78 (0.17)	0.42 (0.05)	2.22 (0.08)
	QUDA	26.51 (0.20)	0.29 (0.07)	25.48 (2.75)	0.82 (0.08)	4.14 (0.12)
	Oracle	21.93 (0.08)	—	—	—	—
	PLR	38.82 (0.33)	9.31 (1.99)	—	1.58 (0.05)	—
500	DSDA	37.10 (0.29)	16.06 (3.02)	—	0.42 (0.05)	—
	sQDA	28.22 (0.41)	24.22 (5.04)	—	0.58 (0.05)	—
	PLR2	35.45 (0.64)	0.34 (0.09)	55.69 (12.67)	0.99 (0.05)	3.05 (0.10)
	IIS-SQDA	26.78 (0.31)	3.22 (1.09)	0.23 (0.05)	0.98 (0.02)	2.65 (0.09)
	QUDA	26.68 (0.27)	0.14 (0.06)	10.96 (1.38)	1.02 (0.08)	4.36 (0.09)
	Oracle	21.81 (0.09)	—	—	—	—

4.2 Real data

In this section, we investigate the performance of QUDA by analyzing two real data sets and compare it to the other classifiers discussed in the simulation study.

Table 2: The means and standard errors (in parentheses) of various performance measures by different classification methods for model 2 based on 100 replications

p	Method	MR (%)	FP.main	FP.inter	FN.main	FN.inter
50	LDA	34.53 (0.19)	—	—	—	—
	QDA	32.09 (0.25)	—	—	—	—
	PLR	31.58 (0.20)	7.51 (0.55)	—	0.07 (0.03)	—
	DSDA	29.89 (0.16)	8.52 (0.86)	—	0.16 (0.04)	—
	sQDA	30.96 (0.90)	27.33 (1.95)	—	0.24 (0.05)	—
	PLR2	5.85 (0.10)	1.14 (0.11)	45.60 (1.08)	0.14 (0.04)	14.27 (0.33)
	IIS-SQDA	5.85 (0.10)	1.14 (0.11)	45.60 (1.08)	0.14 (0.04)	14.27 (0.32)
	QUDA	1.84 (0.08)	4.12 (0.49)	110.10 (10.54)	0.28 (0.05)	1.28 (0.22)
	Oracle	0.65 (0.02)	—	—	—	—
200	PLR	33.34 (0.21)	10.79 (0.70)	—	0.16 (0.04)	—
	DSDA	30.37 (0.23)	11.91 (2.19)	—	0.29 (0.05)	—
	sQDA	33.28 (0.58)	101.75 (7.72)	—	0.27 (0.05)	—
	PLR2	1.73 (0.06)	0.01 (0.01)	12.68 (0.56)	1.08 (0.05)	119.95 (0.52)
	IIS-SQDA	3.98 (0.10)	2.10 (0.15)	15.76 (0.60)	0.11 (0.04)	153.47 (0.31)
	QUDA	0.39 (0.18)	9.03 (2.12)	724.35 (19.52)	0.21 (0.04)	6.05 (0.35)
	Oracle	0 (0)	—	—	—	—
	Oracle	0 (0)	—	—	—	—
500	PLR	34.04 (0.24)	11.17 (1.02)	—	0.30 (0.05)	—
	DSDA	30.99 (0.22)	14.61 (2.64)	—	0.44 (0.05)	—
	sQDA	36.92 (0.64)	243.9 (21.2)	—	0.35 (0.05)	—
	PLR2	1.68 (0.06)	0 (0)	5.52 (0.33)	1.19 (0.05)	401.47 (0.59)
	IIS-SQDA	4.12 (0.09)	2.74 (0.25)	8.02 (0.43)	0.12 (0.04)	451.13 (0.29)
	QUDA	0.16 (0.22)	24.33 (2.18)	4.81e3 (290.1)	0.52 (0.05)	58.09 (1.10)
	Oracle	0 (0)	—	—	—	—
	Oracle	0 (0)	—	—	—	—

Table 3: The means and standard errors (in parentheses) of various performance measures by different classification methods for model 3 based on 100 replications

p	Method	MR (%)	FP.main	FP.inter	FN.main	FN.inter
50	LDA	38.82 (0.19)	—	—	—	—
	QDA	47.57 (0.11)	—	—	—	—
	PLR	36.06 (0.23)	7.73 (0.58)	—	0.14 (0.03)	—
	DSDA	34.82 (0.24)	9.54 (1.09)	—	0.26 (0.04)	—
	sQDA	41.52 (0.51)	14.89 (1.69)	—	0.44 (0.05)	—
	PLR2	37.36 (0.34)	0.60 (0.10)	31.10 (3.21)	0.39 (0.06)	0 (0)
	IIS-SQDA	35.10 (0.22)	5.25 (0.46)	10.85 (0.96)	0.06 (0.02)	0 (0)
	QUDA	34.99 (0.58)	0.82 (0.20)	23.84 (6.69)	0.35 (0.07)	0 (0)
	Oracle	31.68 (0.10)	—	—	—	—
200	PLR	38.50 (0.31)	12.90 (1.08)	—	0.23 (0.04)	—
	DSDA	36.27 (0.28)	14.81 (2.26)	—	0.41 (0.05)	—
	sQDA	43.82 (0.53)	53.18 (6.74)	—	0.52 (0.05)	—
	PLR2	40.31 (0.45)	0.15 (0.05)	40.38 (5.05)	0.74 (0.06)	0 (0)
	IIS-SQDA	36.32 (0.25)	25.39 (0.66)	6.03 (0.50)	0 (0)	0 (0)
	QUDA	36.55 (0.74)	1.70 (1.38)	37.15 (16.39)	0.89 (0.09)	0 (0)
	Oracle	31.54 (0.10)	—	—	—	—
	Oracle	31.54 (0.10)	—	—	—	—
500	PLR	39.98 (0.32)	14.79 (1.41)	—	0.40 (0.05)	—
	DSDA	37.07 (0.29)	19.49 (3.65)	—	0.59 (0.05)	—
	sQDA	46.00 (0.48)	130.91 (18.08)	—	0.57 (0.05)	—
	PLR2	42.23 (0.53)	0.03 (0.02)	36.6 (4.32)	1.07 (0.06)	0 (0)
	IIS-SQDA	37.45 (0.26)	14.53 (1.38)	3.70 (0.32)	0.07 (0.26)	0 (0)
	QUDA	37.95 (0.76)	0.2 (0.06)	57.49 (14.74)	1.05 (0.09)	0 (0)
	Oracle	31.85 (0.12)	—	—	—	—
	Oracle	31.85 (0.12)	—	—	—	—

Table 4: The means and standard errors (in parentheses) of various performance measures by different classification methods for model 4 based on 100 replications

p	Method	MR (%)	FP.main	FP.inter	FN.main	FN.inter
50	LDA	35.58 (0.20)	—	—	—	—
	QDA	35.40 (0.20)	—	—	—	—
	PLR	32.42 (0.23)	8.03 (0.57)	—	0.03 (0.01)	—
	DSDA	31.39 (0.21)	11.02 (1.13)	—	0.09 (0.03)	—
	sQDA	40.90 (0.46)	18.36 (1.93)	—	0.45 (0.05)	—
	PLR2	22.42 (0.21)	1.88 (0.16)	81.56 (2.26)	0.06 (0.03)	123.72 (0.36)
	IIS-SQDA	21.77 (0.20)	3.42 (0.21)	58.92 (1.86)	0 (0)	125.73 (0.32)
	QUDA	16.91 (0.27)	0.55 (0.14)	194.98 (11.31)	0.61 (0.08)	106.51 (0.83)
	Oracle	3.22 (0.04)	—	—	—	—
200	PLR	34.93 (0.28)	12.71 (0.88)	—	0.10 (0.03)	—
	DSDA	32.64 (0.26)	15.63 (2.14)	—	0.21 (0.04)	—
	sQDA	41.68 (0.54)	64.88 (7.33)	—	0.46 (0.05)	—
	PLR2	21.82 (0.20)	0.30 (0.05)	107.80 (2.32)	0.40 (0.05)	559.23 (0.63)
	IIS-SQDA	20.15 (0.19)	6.11 (0.31)	70.76 (1.76)	0 (0)	563.33 (0.38)
	QUDA	9.59 (0.19)	0.31 (0.08)	297.38 (25.33)	0.82 (0.09)	498.61 (1.49)
	Oracle	0.28 (0.02)	—	—	—	—
500	PLR	37.19 (0.32)	15.68 (1.27)	—	0.32 (0.04)	—
	DSDA	33.83 (0.30)	22.90 (3.54)	—	0.45 (0.05)	—
	sQDA	43.39 (0.48)	193.04 (20.32)	—	0.46 (0.05)	—
	PLR2	23.06 (0.23)	0.05 (0.02)	114.94 (2.34)	0.79 (0.05)	1455 (0.65)
	IIS-SQDA	19.07 (0.17)	12.86 (0.42)	57.44 (1.41)	0 (0)	1459 (0.34)
	QUDA	4.18 (0.13)	0.20 (0.04)	298.24 (20.8)	0.42 (0.07)	1315 (2.41)
	Oracle	0 (0)	—	—	—	—

Table 5: The means and standard errors (in parentheses) of various performance measures by different classification methods for model 5 based on 100 replications

p	Method	MR (%)	FP.main	FP.inter	FN.main	FN.inter
50	LDA	39.21 (0.20)	—	—	—	—
	QDA	46.41 (0.17)	—	—	—	—
	PLR	35.76 (0.26)	6.08 (0.43)	—	0.01 (0.01)	—
	DSDA	33.73 (0.25)	8.08 (0.99)	—	0.14 (0.04)	—
	sQDA	36.76 (0.27)	9.37 (1.57)	—	0.06 (0.02)	—
	PLR2	36.62 (0.39)	1.04 (0.13)	45.83 (3.99)	0.05 (0.02)	63.69 (0.39)
	IIS-SQDA	35.56 (0.29)	8.77 (0.50)	14.85 (0.83)	0 (0)	61.18 (0.26)
	QUDA	34.32 (0.53)	0.52 (0.12)	39.76 (6.47)	0.58 (0.08)	59.76 (0.64)
	Oracle	32.36 (0.25)	—	—	—	—
200	PLR	37.73 (0.34)	9.68 (0.89)	—	0.40 (0.03)	—
	DSDA	34.58 (0.35)	10.87 (2.44)	—	0.11 (0.03)	—
	sQDA	26.11 (0.27)	18.35 (4.79)	—	0.21 (0.04)	—
	PLR2	37.40 (0.44)	0.32 (0.06)	66.44 (5.47)	0.31 (0.06)	194.46 (0.35)
	IIS-SQDA	33.22 (0.28)	19.87 (0.93)	6.16 (0.41)	0 (0)	191.37 (0.10)
	QUDA	29.35 (0.41)	0.10 (0.05)	164.24 (73.3)	1.27 (0.07)	175.8 (0.96)
	Oracle	20.09 (0.27)	—	—	—	—
500	PLR	39.13 (0.33)	14.39 (1.29)	—	0.08 (0.03)	—
	DSDA	34.76 (0.25)	9.44 (1.77)	—	0.16 (0.04)	—
	sQDA	10.17 (0.16)	22.32 (6.88)	—	0.24 (0.05)	—
	PLR2	37.44 (0.52)	0.16 (0.05)	90.78 (6.06)	0.43 (0.06)	493.48 (0.41)
	IIS-SQDA	26.57 (0.23)	19.14 (0.57)	62.00 (1.56)	0 (0)	475.49 (0.20)
	QUDA	23.75 (0.49)	4.03 (2.91)	507.92 (225.36)	1.59 (0.06)	459.96 (1.96)
	Oracle	4.16 (0.08)	—	—	—	—

Quora answer classifier. This is a data challenge available at http://www.quora.com/challenges#answer_classifier. The training data set contains 4,500 answers from QUORA which have been annotated with either "good" or "bad". For each answer, 21 features (20 of which are effective) were extracted from the original sentences. The goal of this challenge is to automatically classify a new answer based on the 20 features. Since the dimension $p = 20$ is relatively small, we can compare QUDA to all the methods discussed in the simulation via 5-fold cross-validation. In particular, we randomly split the data into five parts, fit a model to the four parts of the data, and report the misclassification error on the part that is left out. The average misclassification errors and the standard errors for various methods are in Table 6. Interestingly, LDA performs much better than QDA, suggesting that if we stop the analysis here, we might simply have preferred to use the linear classifier LDA. However, the story becomes different if sparse models are considered. In particular, PLR, PLR2, IIS-SQDA and QUDA all outperform the non-sparse models significantly with QUDA performing the best.

Table 6: Misclassification rate (%) for the Quora answer data under 5-fold cross-validation

Method	mean	standard error
LDA	18.84	0.50
QDA	30.33	0.72
PLR	17.89	0.60
DSDA	19.11	0.56
sQDA	29.59	1.57
PLR2	17.56	0.71
IIS-SQDA	17.33	0.48
QUDA	16.44	0.45

Prostate cancer. Taken from <ftp://stat.ethz.ch/Manuscripts/dettling/prostate.rda>, this data contains genetic expression levels for $N = 6033$ genes of 102 individuals. The first 50 are normal control subjects while the rest are prostate cancer patients. More details of the data can be found in Singh, et al. (2002), Dettling (2004) and Efron (2010). The goal is to identify genes that are linked with prostate cancer and predict potential patients and the difficulty of this task lies in the interactions among genes. The existence of interactions

can often complicate the analysis and produce unreliable inference if they are ignored. For example, Figure 1 displays the pair of 118th and 182th gene. We can see the marginal distributions of each gene does not differ too much between the patients and the normal subjects (the middle and the right panels), suggesting that their main effects may not be important for distinguishing the two classes. In the left panel of Figure 1, however, we can identify some joint pattern that distinguishes the two groups. It can be seen that most patients are allocated in the red triangle while most normal subjects are within the blue triangle, indicating the existence of some interaction effect that might be useful for classification.




Figure 1: The plot for the gene 118 and gene 182. Left: joint scatter plot; Middle: marginal density of gene 118; Right: marginal density of gene 182.

For this data, we follow the same method in Cai and Liu (2011), retaining only the top 200 or 500 genes with the largest absolute values of the two sample t statistics. The average misclassification errors and the standard errors using 5-fold cross-validation for various methods are reported in Table 7. Note that since $p \gg n$, LDA and QDA were excluded. We can see again that QUDA outperforms all the other methods by a large margin, regardless of the number of the genes that were used for analysis.

Table 7: Misclassification rate (%) for the prostate cancer data under 5-fold cross-validation

Method	$p = 200$		$p = 500$	
	mean	std error	mean	std error
PLR	11.00	2.45	13.00	4.06
DSDA	5.00	3.32	11.00	2.92
sQDA	3.00	1.22	2.00	1.22
PLR2	13.00	4.47	23.00	3.39
IIS-SQDA	11.00	2.92	15.00	2.74
QUDA	1.00	1.00	2.00	1.22

5 Conclusion

We have proposed a novel method named QUDA for high-dimensional quadratic discriminant analysis. This is the first method aiming at directly estimating the quantities in the QDA discriminant function. The proposed framework is simple, fast to implement and enjoys excellent theoretical properties. We have demonstrated via extensive simulation and two data analyses that QUDA performs competitively under various circumstances.

We conclude by identifying three directions for future research. First, as the discussion of the paper is focused on binary problems, it will be interesting to extend it to multi-class problems. Second, it is also interesting to see whether our theoretical results are optimal and in what sense. Finally, the proposed framework is extremely flexible. As a concrete example, if Ω is a two block sparse matrix after permutation as in Fan et al. (2015b), we can change the penalty $\|\Omega\|_1$ in (2) to one that encourages row sparsity, for example to $\|\Omega\|_{1,2} = \sum_{j=1}^p \|\Omega_{i,:}\|_2$ which is the sum of the ℓ_2 norms of the rows. It will be interesting to see how well this procedure compares with IIS-SQDA in Fan et al. (2015b). This topic is beyond the scope of the current paper and will be pursued elsewhere.

References

- Anderson, T.W. (2003). *An Introduction to Multivariate Statistical Analysis*, 3rd ed. Wiley Series in Probability and Statistics. Wiley, New York.
- Bien, J., Taylor, J., and Tibshirani, R. (2013). A lasso for hierarchical interactions. *The Annals of Statistics*, 41, 1111-1141.

- Bickel, P. and Levina, E. (2004). Some theory for fisher’s linear discriminant function, ‘naive bayes’, and some alternatives when there are many more variables than observations. *Bernoulli*, 10, 989-1010.
- Bickel, P. J., Levina, E. (2008). Covariance regularization by thresholding. *The Annals of Statistics* 36, 199-227.
- Boyd, S., Parikh, N., Chu, E., Peleato, B., and Eckstein, J. (2011). Distributed optimization and statistical learning via the alternating direction method of multipliers. *Foundations and Trends in Machine Learning*, 3, 1-122.
- Cai, T. and Liu, W. (2011). A direct estimation approach to sparse linear discriminant analysis. *Journal of the American Statistical Association*, 106, 1566-1577.
- Detting, M. (2004). BagBoosting for tumor classification with gene expression data. *Bioinformatics* 20, 3583-3593.
- Donoho, D., and Huo, X. (2001). Uncertainty principles and ideal atomic decomposition. *Information Theory, IEEE Transactions*, 47, 2845-2862.
- Efron, B. (2010). Large-scale inference: empirical Bayes methods for estimation, testing, and prediction. Vol. 1. Cambridge University Press.
- Fan, J., Feng, Y., and Tong, X. (2012). A ROAD to classification in high dimensional space. *Journal of the Royal Statistical Society Series B*, 74, 745-771.
- Fan, J., Ke, T., Liu, H., and Xia, L. (2015). QUADRO: A supervised dimension reduction method via Rayleigh quotient optimization. *The Annals of Statistics*, 43, 1493-1534.
- Fan, Y., Kong, Y., Li, D., and Zheng, Z. (2015). Innovated interaction screening for high-dimensional nonlinear classification. *The Annals of Statistics*, 43, 1243-1272.
- Friedman, J., Hastie, T., and Tibshirani, R. (2010). Regularization paths for generalized linear models via coordinate descent. *Journal of statistical software*, 33, 1.
- Ghadimi, E., Teixeira, A., Shames, I., and Johansson, M. (2015). Optimal parameter selection for the alternating direction method of multipliers (ADMM): quadratic problems. *Automatic Control, IEEE Transactions on*, 60(3), 644-658.
- Hao, N., Feng, Y., and Zhang, H. H. (2014). Variable selection for high dimensional quadratic regressions via regularization. <http://arxiv.org/abs/1501.00049>.
- Hao, N. and Zhang, H. H. (2014). Interaction screening for ultra-high dimensional data. *Journal of the American Statistical Association*, 109, 1285–1301.
- Hao, N. and Zhang, H.H. (2015). A note on high dimensional linear regression with interactions. <http://arxiv.org/abs/1412.7138>.
- Hastie, T. J., Tibshirani, R. J., and Friedman, J. F. (2009). The elements of statistical learning: data mining, inference, and prediction. Springer.

- Jiang, B., Chen, Z., and Leng, C. (2015). Dynamic linear discriminant analysis for high-dimensional data. Manuscript.
- Leng, C. (2008). Sparse optimal scoring for multiclass cancer diagnosis and biomarker detection using microarray data. *Computational Biology and Chemistry*, 32, 417-425.
- Li, Q. and Shao, J. (2015). Sparse quadratic discriminant analysis for high dimensional data. *Statistica Sinica*, 25, 457-473.
- Liu, H., Lafferty, J., and Wasserman, L. (2009). The nonparanormal: Semiparametric estimation of high dimensional undirected graphs. *Journal of Machine Learning Research*, 10, 2295-2328.
- Mai, Q., Yang, Y., and Zou, H. (2015). Multiclass Sparse Discriminant Analysis. Manuscript.
- Mai, Q. and Zou, H. (2013). A note on the equivalence of three sparse linear discriminant methods. *Technometrics*, 55, 243-246.
- Mai, Q., Zou, H., and Yuan, M. (2012). A direct approach to sparse discriminant analysis in ultra-high dimensions. *Biometrika*, 99, 29-42.
- Pan, R., Wang, H., and Li, R. (2015). Ultrahigh dimensional multi-class linear discriminant analysis by pairwise sure independence screening. *Journal of the American Statistical Association*, to appear.
- Shao, J., Wang, Y., Deng, X., and Wang, S. (2011). Sparse linear discriminant analysis by thresholding for high dimensional data. *The Annals of statistics*, 39, 1241-1265.
- Ravikumar, P., Wainwright, M., Raskutti, G., Yu, B. (2011). High-dimensional covariance estimation by minimizing l1-penalized log-determinant divergence. *Electronic Journal of Statistics*, 5, 935-980.
- Singh, D., Febbo, P. G., Ross, K., Jackson, D. G., Manola, J., Ladd, C. et al. (2002). Gene expression correlates of clinical prostate cancer behavior. *Cancer cell*, 1, 203-209.
- Sun, J. and Zhao, H. (2015). The application of sparse estimation of covariance matrix to quadratic discriminant analysis. *BMC Bioinformatics*, 16, 48.
- Witten, D. M. and Tibshirani, R. (2011). Penalized classification using Fisher's linear discriminant. *Journal of the Royal Statistical Society Series B*, 73, 753-772.
- Van der Vaart, A. W. (2000). *Asymptotic statistics*, Vol. 3. Cambridge university press.
- Zhang, T. and Zou, H. (2014). Sparse precision matrix estimation via lasso penalized D-trace loss. *Biometrika*, 101, 103-120.
- Zhao, J. and Leng, C. (2015). An analysis of penalised interaction models. *Bernoulli*, to appear.

- Zhao, P. and Yu, B. (2006). On model selection consistency of Lasso. *The Journal of Machine Learning Research*, 7, 2541-2563.
- Zhao, S. D., Cai, T. T., and Li, H. (2014). Direct estimation of differential networks. *Biometrika*, 101, 253–268.

Supplementary Materials: technical details

Proofs of Theorem 1

We first introduce some technical lemmas and the the proof of Theorem 1 will be given after these lemmas.

Lemma 1. *Suppose $\lambda_{\max}(\Sigma_k) < \epsilon_0 < \infty$ for $k = 1, 2$. There exist constants $C_0, C_1, C_2 > 0$ depending on ϵ_0 only, such that for any $|v| \leq C_0$,*

$$P(|\hat{\sigma}_{kij} - \sigma_{kij}| > v) \leq C_1 \exp(-C_2(n_k - 1)v^2) \leq C_1 \exp(-C_2nv^2).$$

Proof. Denote $X = (X_1, \dots, X_{n_1})^T$. Let Λ be an orthogonal matrix with the last row being $(n^{-1/2}, \dots, n^{-1/2})$ and define $Z = (z_1, \dots, z_n) = \Lambda X$. We then have $z_1, \dots, z_{n-1} \sim N(\mathbf{0}, \Sigma)$ and they are independent to each other. Note that $n_1 \hat{\Sigma}_1 = X^T(I_p - \mathbf{1}\mathbf{1}^T)X = Z^T\Lambda(I_p - \mathbf{1}\mathbf{1}^T)\Lambda^T Z = \sum_{i=1}^{n-1} z_i z_i^T$. This together with Lemma A.3 of Bickel and Levina (2008) prove Lemma 1. $\square$

Remark. Denote $\sigma^2 = \max\{\sigma_{1ii}, \sigma_{2ii}, i = 1, \dots, p\}$. From Lemma 1 of Ravikumar et al. (2011) we can see that Lemma 1 is true for $C_1 = 4$, $C_2 = [128(1 + 4\sigma^2)^2\sigma^4]^{-1}$ and $v = 8(1 + 4\sigma^2)\sigma^2$.

Lemma 2. *Assume that,*

$$B_{\Gamma, \Gamma^T} < \frac{1}{3(d^2\epsilon_1\epsilon_2 + Bd^2(\epsilon_1 + \epsilon_2))}. \quad (9)$$

Let $R(\Delta_\Gamma) = (\Gamma_{S,S} + \Delta_\Gamma)^{-1} - \Gamma_{S,S}^{-1} + \Gamma_{S,S}^{-1}(\Delta_\Gamma)_{S,S}\Gamma_{S,S}^{-1}$. We then have

$$\|R(\Delta_\Gamma)\|_\infty \leq 3\|(\Delta_\Gamma)_{S,S}\|_\infty\|(\Delta_\Gamma^T)_{S,S}\|_{1,\infty}B_\Gamma B_{\Gamma^T}^2, \quad (10)$$

and

$$\|R(\Delta_\Gamma)\|_{1,\infty} \leq 3\|(\Delta_\Gamma)_{S,S}\|_{1,\infty}\|(\Delta_{\Gamma^T})_{S,S}\|_{1,\infty}B_\Gamma B_{\Gamma^T}^2. \quad (11)$$

Moreover, we also have

$$\|\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}\|_\infty \leq 3d^2(\epsilon^2 + 2B\epsilon)^2 B_\Gamma B_{\Gamma^T}^2 + (\epsilon^2 + 2B\epsilon)B_\Gamma^2, \quad (12)$$

$$\|\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}\|_{1,\infty} \leq 3d^4(\epsilon^2 + 2B\epsilon)^2 B_\Gamma B_{\Gamma^T}^2 + d^2(\epsilon^2 + 2B\epsilon)B_\Gamma^2. \quad (13)$$

Proof. Note that

$$\Delta_\Gamma = \Delta_2 \otimes \Delta_1 + \Delta_2 \otimes \Sigma_1 + \Sigma_2 \otimes \Delta_1.$$

Consequently by (9) we have $\|\Gamma_{S,S}^{-1}\|_{1,\infty}\|(\Delta_\Gamma)_{S,S}\|_{1,\infty} \leq 1/3$. (10) and (11) can then be proved using the same arguments as in Appendix B of Ravikumar et al. (2011). Note that $\|(\Delta_\Gamma)_{S,S}\|_\infty \leq \epsilon^2 + 2B\epsilon$ and $\max\{\|(\Delta_\Gamma)_{S,S}\|_{1,\infty}, \|(\Delta_{\Gamma^T})_{S,S}\|_{1,\infty}\} \leq d^2(\epsilon^2 + 2B\epsilon)$, we have

$$\begin{aligned} \|\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}\|_\infty &\leq \|R(\Delta_\Gamma)\|_\infty + \|\Gamma_{S,S}^{-1}(\Delta_\Gamma)_{S,S}\Gamma_{S,S}^{-1}\|_\infty \\ &\leq 3d^2(\epsilon^2 + 2B\epsilon)^2 B_\Gamma B_{\Gamma^T}^2 + \|(\Delta_\Gamma)_{S,S}\|_\infty \|\Gamma_{S,S}^{-1}\|_{1,\infty}^2 \\ &\leq 3d^2(\epsilon^2 + 2B\epsilon)^2 B_\Gamma B_{\Gamma^T}^2 + (\epsilon^2 + 2B\epsilon)B_\Gamma^2. \end{aligned}$$

This proves (12). (13) can be proved similarly. $\square$

Lemma 3. Assume that (9) and the following assumptions hold: $\alpha > 0$, $\epsilon < \min\left\{B, \frac{\alpha\lambda}{2(2-\alpha)}\right\}$

and

$$3d^2\epsilon BB_{\Gamma,\Gamma^T}[1 + (B_\Sigma^2 + 3d^2\epsilon BB_{\Gamma,\Gamma^T})(9d^2\epsilon BB_{\Gamma,\Gamma^T} + 1)B_{\Gamma,\Gamma^T}] \leq C_\alpha \alpha \min\{\lambda, 1\}, \quad (14)$$

where $C_\alpha = \frac{\alpha\lambda + 2\epsilon\alpha - 4\epsilon}{2B\alpha\lambda + \alpha\lambda + 2\epsilon\alpha}$. We have:

$$(i) \text{ } \text{vec}(\hat{\Omega})_S = 0.$$

$$(ii) \|\hat{\Omega} - \Omega\|_\infty < 2\lambda B_{\Gamma, \Gamma^T} + 9d^2\epsilon B B_{\Gamma, \Gamma^T}^2 (3d^2\epsilon B B_{\Gamma, \Gamma^T} + 1)(2B + 2\lambda).$$

Proof. (i) Suppose $\tilde{\Omega}$ is the solution of:

$$\tilde{\Omega} = \min_{\Omega \in R^{p \times p}, \Omega_{S^c} = 0} \frac{1}{2} \text{Tr} \left(\Omega^T \hat{\Sigma}_1 \Omega \hat{\Sigma}_2 \right) - \text{Tr} \left(\Omega (\hat{\Sigma}_1 - \hat{\Sigma}_2) \right) + \lambda \|\Omega\|_1. \quad (15)$$

We prove Lemma 3 (i) by showing that $\hat{\Omega} = \tilde{\Omega}$. Due to the convexity of (3) in the main paper, we only need to show that the derivative of (3) is zero at $\tilde{\Omega}$. Equivalently, we need to show that for any $1 \leq i, j \leq p$ we have,

$$|\hat{\Sigma}_1 \tilde{\Omega} \hat{\Sigma}_2 - (\hat{\Sigma}_1 - \hat{\Sigma}_2)|_{i,j} \leq \lambda. \quad (16)$$

By taking the first derivative of (15) we obtain,

$$\{\hat{\Sigma}_1 \tilde{\Omega} \hat{\Sigma}_2 - (\hat{\Sigma}_1 - \hat{\Sigma}_2) + \lambda Z\}_S = \mathbf{0}, \quad (17)$$

where $Z = (Z_{ij})_{1 \leq i, j \leq p}$ with $Z_{ij} = 0$ for $(i, j) \in S^c$, $Z_{ij} = \text{sign}(\tilde{\Omega}_{ij})$ for $(i, j) \in S$ and $\tilde{\Omega}_{ij} \neq 0$, $Z_{ij} \in [-1, 1]$ for $(i, j) \in S$ and $\tilde{\Omega}_{ij} = 0$. Therefore (16) is true for any $(i, j) \in S$. Using the vector operator, (17) becomes

$$\{(\hat{\Sigma}_2 \otimes \hat{\Sigma}_1) \text{vec}(\tilde{\Omega}) - \text{vec}(\hat{\Sigma}_1 - \hat{\Sigma}_2) + \lambda \text{vec}(Z)\}_S = \mathbf{0}.$$

Equivalently we have

$$\text{vec}(\tilde{\Omega})_S = \hat{\Gamma}_{S, S}^{-1} [\text{vec}(\hat{\Sigma}_1 - \hat{\Sigma}_2)_S - \lambda \text{vec}(Z)_S]. \quad (18)$$

Note that the left hand side of (16) equals $|\hat{\Sigma}_1 \tilde{\Omega} \hat{\Sigma}_2 - \Sigma_1 + \Sigma_2 - (\Delta_1 - \Delta_2)|_{i,j}$. Using the vector operator and the fact that $\tilde{\Omega}_{S^c} = \mathbf{0}$, to show that (16) is true for any $e \in S^c$, we only

need to show that

$$|\hat{\Gamma}_{e,S} \text{vec}(\tilde{\Omega})_S - \Gamma_{e,S} \text{vec}(\Omega)_S - \text{vec}(\Delta_1 - \Delta_2)_e| \leq \lambda. \quad (19)$$

Here we have use the fact that $\Gamma_{e,S} \text{vec}(\Omega)_S = \text{vec}(\Sigma_1 - \Sigma_2)_e$. By (18) and the fact that $\text{vec}(\Omega)_S = \Gamma_{S,S}^{-1} \text{vec}(\Sigma_1 - \Sigma_2)_S$ we have,

$$\begin{aligned} & |\hat{\Gamma}_{e,S} \text{vec}(\tilde{\Omega})_S - \Gamma_{e,S} \text{vec}(\Omega)_S - \text{vec}(\Delta_1 - \Delta_2)_e| \\ &= |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1} [\text{vec}(\hat{\Sigma}_1 - \hat{\Sigma}_2)_S - \lambda \text{vec}(Z)_S] - \Gamma_{e,S} \Gamma_{S,S}^{-1} \text{vec}(\Sigma_1 - \Sigma_2)_S - \text{vec}(\Delta_1 - \Delta_2)_e| \\ &\leq |[\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1} - \Gamma_{e,S} \Gamma_{S,S}^{-1}] \text{vec}(\Sigma_1 - \Sigma_2)_S| + |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1} \text{vec}(\hat{\Sigma}_1 - \hat{\Sigma}_2 - \Sigma_1 + \Sigma_2)_S| \\ &\quad + \|\Delta_1 - \Delta_2\|_\infty + \lambda |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1}|_1 \\ &\leq 2B |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1} - \Gamma_{e,S} \Gamma_{S,S}^{-1}|_1 + (\epsilon_1 + \epsilon_2 + \lambda) |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1}|_1 + \epsilon_1 + \epsilon_2. \end{aligned}$$

Consequently, (19) is true if

$$\begin{aligned} & \max_{e \in S^c} 2B |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1} - \Gamma_{e,S} \Gamma_{S,S}^{-1}|_1 + 2\epsilon(1 + |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1}|_1) \leq (1 - C_\alpha) \alpha \lambda, \\ & \max_{e \in S^c} |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1}|_1 \leq 1 - (1 - C_\alpha) \alpha. \end{aligned} \quad (20)$$

Next we finish this proof by showing that (20) is true under the assumptions of this lemma.

By Lemma 2 we have for any $e \in S^c$,

$$\begin{aligned} & |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1} - \Gamma_{e,S} \Gamma_{S,S}^{-1}|_1 \\ &\leq |(\hat{\Gamma}_{e,S} - \Gamma_{e,S}) \Gamma_{S,S}^{-1}|_1 + |\Gamma_{e,S} (\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1})|_1 + |(\hat{\Gamma}_{e,S} - \Gamma_{e,S}) (\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1})|_1 \\ &\leq |\hat{\Gamma}_{e,S} - \Gamma_{e,S}|_1 \|\Gamma_{S,S}^{-1}\|_{1,\infty} + \|\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}\|_{1,\infty} B_\Sigma^2 + |\hat{\Gamma}_{e,S} - \Gamma_{e,S}|_1 \|\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}\|_{1,\infty} \\ &\leq d^2(\epsilon^2 + 2B\epsilon) B_{\Gamma,\Gamma^T} + d^2[B_\Sigma^2 + d^2(\epsilon^2 + 2B\epsilon)] [3d^2(\epsilon^2 + 2B\epsilon) B_{\Gamma,\Gamma^T} + 1] (\epsilon^2 + 2B\epsilon) B_{\Gamma,\Gamma^T}^2 \\ &\leq 3d^2 \epsilon B B_{\Gamma,\Gamma^T} [1 + (B_\Sigma^2 + 3d^2 \epsilon B B_{\Gamma,\Gamma^T}) (9d^2 \epsilon B B_{\Gamma,\Gamma^T} + 1) B_{\Gamma,\Gamma^T}] \\ &\leq C_\alpha \alpha \min\{\lambda, 1\}. \end{aligned}$$

Consequently, $\max_{e \in S^c} |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1}|_1 \leq C_\alpha \alpha \min\{\lambda, 1\} + (1 - \alpha) \leq 1 - (1 - C_\alpha)\alpha$, and

$$\begin{aligned} & \max_{e \in S^c} 2B |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1} - \Gamma_{e,S} \Gamma_{S,S}^{-1}|_1 + 2\epsilon(1 + |\hat{\Gamma}_{e,S} \hat{\Gamma}_{S,S}^{-1}|_1) \\ & \leq 2BC_\alpha \alpha \lambda + 2\epsilon[2 - (1 - C_\alpha)\alpha] \\ & = (1 - C_\alpha)\alpha \lambda. \end{aligned}$$

(ii) From (i) we have $\hat{\Omega} = \tilde{\Omega}$. By (18) and the fact that $\text{vec}(\Omega)_S = \Gamma_{S,S}^{-1} \text{vec}(\Sigma_1 - \Sigma_2)_S$, we have

$$\begin{aligned} \|\hat{\Omega} - \Omega\|_\infty &= |\text{vec}(\tilde{\Omega}) - \text{vec}(\Omega)|_\infty \\ &= |\hat{\Gamma}_{S,S}^{-1} [\text{vec}(\hat{\Sigma}_1 - \hat{\Sigma}_2)_S - \lambda \text{vec}(Z)_S] - \Gamma_{S,S}^{-1} \text{vec}(\Sigma_1 - \Sigma_2)_S|_\infty \\ &\leq \lambda \|\hat{\Gamma}_{S,S}^{-1}\|_{1,\infty} + |\hat{\Gamma}_{S,S}^{-1} \text{vec}(\Delta_1 - \Delta_2)_S|_\infty + |(\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}) \text{vec}(\Sigma_1 - \Sigma_2)|_\infty \\ &\leq (\lambda + 2\epsilon) \|\hat{\Gamma}_{S,S}^{-1}\|_{1,\infty} + 2B \|\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}\|_{1,\infty} \\ &\leq (\lambda + 2\epsilon) \|\Gamma_{S,S}^{-1}\|_{1,\infty} + (2B + \lambda + 2\epsilon) \|\hat{\Gamma}_{S,S}^{-1} - \Gamma_{S,S}^{-1}\|_{1,\infty}. \end{aligned}$$

By (13) and the assumption that $2\epsilon < \alpha\lambda/(2 - \alpha) < \alpha\lambda < \lambda$ we immediately have

$$\begin{aligned} \|\hat{\Omega} - \Omega\|_\infty &\leq (\lambda + 2\epsilon) B_\Gamma + 3[d^4(\epsilon^2 + 2B\epsilon)^2 B_\Gamma B_{\Gamma^T}^2 + d^2(\epsilon^2 + 2B\epsilon) B_\Gamma^2](2B + \lambda + 2\epsilon) \\ &< 2\lambda B_{\Gamma, \Gamma^T} + 9d^2 \epsilon B B_{\Gamma, \Gamma^T}^2 (3d^2 \epsilon B B_{\Gamma, \Gamma^T} + 1)(2B + 2\lambda). \end{aligned}$$

□

Proof of Theorem 1

From Lemma 1 we have that with probability greater than $1 - p^{2-c}$, $\epsilon \leq \{(c \log p + \log C_1)/C_2 n\}^{1/2}$.

With an abuse of notation, we denote $\epsilon = \{(c \log p + \log C_1)/C_2 n\}^{1/2}$. We firstly verify that the assumptions in Lemmas 2 and 3 are true for the given λ , n and ϵ .

(i) By noticing that $\frac{8\epsilon}{\alpha} < \lambda$, $\frac{\alpha}{8} < \frac{\alpha}{4(2-\alpha)}$ and $n > (C_2 B^2)^{-1}(c \log p + \log C_1)$ we immediately

have

$$\epsilon < \min \left\{ B, \frac{\alpha\lambda}{4(2-\alpha)} \right\} < \min \left\{ B, \frac{\alpha\lambda}{2(2-\alpha)} \right\}.$$

(ii) $n > 81B^2d^4B_{\Gamma,\Gamma^T}^2(c\log p + \log C_1)/C_2$ implies $B_{\Gamma,\Gamma^T} < \frac{1}{9d^2B\epsilon}$. Together with $\epsilon < B$ from (i) we can see that Assumption (9) holds.

(iii) Since $\epsilon < \frac{\alpha\lambda}{2(2-\alpha)}$, we have

$$C_\alpha > \frac{\alpha\lambda - 4\epsilon}{2B\alpha\lambda + \alpha\lambda} > \frac{\alpha\lambda - 2\alpha\lambda/(2-\alpha)}{2B\alpha\lambda + \alpha\lambda} = \frac{1-\alpha}{(2-\alpha)(2B+1)}.$$

Together with $B_{\Gamma,\Gamma^T} < \frac{1}{9d^2B\epsilon}$, we have

$$\begin{aligned} & 3d^2\epsilon BB_{\Gamma,\Gamma^T}[1 + (B_\Sigma^2 + 3d^2\epsilon BB_{\Gamma,\Gamma^T})(9d^2\epsilon BB_{\Gamma,\Gamma^T} + 1)B_{\Gamma,\Gamma^T}] \\ & < 3d^2\epsilon BB_{\Gamma,\Gamma^T}[1 + 2(B_\Sigma^2 + 1/3)B_{\Gamma,\Gamma^T}] \\ & \leq \frac{1-\alpha}{(2-\alpha)(2B+1)}\lambda \\ & < C_\alpha\lambda. \end{aligned}$$

(14) is then true since $n > 9(C_2\alpha)^{-1}B^2B_{\Gamma,\Gamma^T}^2[1 + 2(B_\Sigma^2 + 1/3)B_{\Gamma,\Gamma^T}]^2(c\log p + \log C_1)$ implies

$$3\epsilon BB_{\Gamma,\Gamma^T}[1 + (B_\Sigma^2 + 3d^2\epsilon BB_{\Gamma,\Gamma^T})(9d^2\epsilon BB_{\Gamma,\Gamma^T} + 1)B_{\Gamma,\Gamma^T}] \leq \alpha.$$

(i), (ii) and (iii) and Lemma 3 imply that

$$\begin{aligned} \|\hat{\Omega} - \Omega\|_\infty & < 2\lambda B_{\Gamma,\Gamma^T} + 9d^2\epsilon BB_{\Gamma,\Gamma^T}^2(3d^2\epsilon BB_{\Gamma,\Gamma^T} + 1)(2B + 2\lambda) \\ & \leq 2\lambda B_{\Gamma,\Gamma^T} + 12d^2\epsilon BB_{\Gamma,\Gamma^T}^2(2B + 2\lambda) \\ & = \frac{14}{3}\lambda B_{\Gamma,\Gamma^T} + 24d^2\epsilon B^2B_{\Gamma,\Gamma^T}^2. \end{aligned}$$

Proofs of Theorem 2

We first introduce some technical lemmas and the the proof of Theorem 2 will be given after these lemmas.

Lemma 4. *Assume that $A_\Sigma d_\delta \epsilon < 1$ we have*

$$\|\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1}\|_{1,\infty} \leq \frac{A_\Sigma^2 d_\delta \epsilon}{1 - A_\Sigma d_\delta \epsilon}.$$

Proof. This lemma can be easily proved using the following observation:

$$\begin{aligned} \|\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1}\|_{1,\infty} &\leq \|\hat{\Sigma}_{D,D}^{-1}\|_{1,\infty} \|\hat{\Sigma}_{D,D} - \Sigma_{D,D}\|_{1,\infty} \|\Sigma_{D,D}^{-1}\|_{1,\infty} \\ &\leq (A_\Sigma + \|\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1}\|_{1,\infty}) d_\delta \epsilon A_\Sigma. \end{aligned}$$

□

Lemma 5.

$$|\hat{\gamma} - \gamma|_\infty \leq 8\epsilon_\mu + 2(\epsilon + B_\Sigma \epsilon_\Omega A_2 + d\epsilon \epsilon_\Omega A_2) |\Omega(\mu_1 - \mu_2)|_1 + 2(B + \epsilon)(A_1 + d\epsilon_\Omega) \epsilon_\mu.$$

Proof.

$$\begin{aligned} |\gamma - \hat{\gamma}|_\infty &\leq 4|\hat{\Delta}_\mu - \Delta_\mu|_\infty + |(\Delta_1 - \Delta_2)\Omega\Delta_\mu|_\infty + |(\hat{\Sigma}_1 - \hat{\Sigma}_2)(\hat{\Omega} - \Omega)\Delta_\mu|_\infty \\ &\quad + |(\hat{\Sigma}_1 - \hat{\Sigma}_2)\hat{\Omega}(\hat{\Delta}_\mu - \Delta_\mu)|_\infty. \end{aligned}$$

Lemma 5 can then be proved using the following facts:

$$|\hat{\Delta}_\mu - \Delta_\mu|_\infty \leq 2\epsilon_\mu$$

$$|\mu_1 - \mu_2|_1 \leq \|(\Omega^{-1})_{\cdot,D}\|_{1,\infty} |\Omega(\mu_1 - \mu_2)|_1 = A_2 |\Omega(\mu_1 - \mu_2)|_1;$$

$$|(\Delta_1 - \Delta_2)\Omega\Delta_\mu|_\infty \leq 2\epsilon|\Omega(\mu_1 - \mu_2)|_1;$$

$$\begin{aligned} |(\hat{\Sigma}_1 - \hat{\Sigma}_2)(\hat{\Omega} - \Omega)\Delta_\mu|_\infty &\leq \|(\Sigma_1 - \Sigma_2)(\hat{\Omega} - \Omega)\|_\infty |\Delta_\mu|_1 + \|(\Delta_1 - \Delta_2)(\hat{\Omega} - \Omega)\|_\infty |\Delta_\mu|_1 \\ &\leq 2B_{\Sigma}\epsilon_\Omega |\mu_1 - \mu_2|_1 + 2d\epsilon_\Omega |\mu_1 - \mu_2|_1; \end{aligned}$$

$$\begin{aligned} |(\hat{\Sigma}_1 - \hat{\Sigma}_2)\hat{\Omega}(\hat{\Delta}_\mu - \Delta_\mu)|_\infty &\leq 2(B + \epsilon)|\hat{\Omega}(\hat{\Delta}_\mu - \Delta_\mu)|_1 \\ &\leq 2(B + \epsilon)[\|\Omega\|_{1,\infty}|\hat{\Delta}_\mu - \Delta_\mu|_\infty + d\epsilon_\Omega|\hat{\Delta}_\mu - \Delta_\mu|_\infty] \\ &\leq 2(B + \epsilon)(A_1 + d\epsilon_\Omega)\epsilon_\mu. \end{aligned}$$

□

Proof of Theorem 2

(i) Suppose $\tilde{\delta}$ is the solution of:

$$\tilde{\delta} = \min_{\delta \in R^p, \delta_{D^c} = 0} \frac{1}{2}\delta^T(\hat{\Sigma}_1 + \hat{\Sigma}_2)\delta - \hat{\gamma}^T\delta + \lambda_\delta\|\delta\|_1,$$

We first show that $\hat{\delta} = \tilde{\delta}$. It suffices to show that for any $e \in D^c$,

$$|2\hat{\Sigma}_{e,D}\tilde{\delta}_D - \hat{\gamma}_e| \leq \lambda_\delta.$$

By the definition of $\tilde{\delta}$ we have

$$\{2\hat{\Sigma}\tilde{\delta} - \hat{\gamma} + \lambda_\delta Z\}_D = \mathbf{0},$$

where $Z = (Z_1, \dots, Z_p)^T$ with $Z_i = 0$ for $i \in D^c$, $Z_i = \text{sign}(\tilde{\delta})$ for $i \in D$ and $\tilde{\delta} \neq 0$, $Z_i \in [-1, 1]$ for $i \in D$ and $\tilde{\delta} = 0$. Consequently, we have $\tilde{\delta}_D = \frac{1}{2}\hat{\Sigma}_{D,D}^{-1}(\hat{\gamma}_D - \lambda_\delta Z_D)$. Together

with the fact that $\Sigma_{e,D}\Sigma_{D,D}^{-1}\gamma_D = 2\Sigma_{e,D}\Sigma_{D,D}^{-1}\Sigma_{D,D}\delta_D = \gamma_e$, we have

$$\begin{aligned}
|2\hat{\Sigma}_{e,D}\tilde{\delta}_D - \hat{\gamma}_e| &= |\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1}(\hat{\gamma}_D - \lambda_\delta Z_D) - \hat{\gamma}_e| \\
&\leq |(\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1} - \Sigma_{e,D}\Sigma_{D,D}^{-1})\gamma_D| + |\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1}(\hat{\gamma}_D - \gamma_D)| \\
&\quad + \lambda_\delta |\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1}|_1 + |\gamma_e - \hat{\gamma}_e| \\
&\leq |(\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1} - \Sigma_{e,D}\Sigma_{D,D}^{-1})|_1 |\gamma_D|_\infty + |\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1}|_1 |(\hat{\gamma}_D - \gamma_D)|_\infty \\
&\quad + \lambda_\delta |\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1}|_1 + |\gamma_e - \hat{\gamma}_e|. \tag{21}
\end{aligned}$$

For simplicity, in the following, inequalities will be derived without mentioning whether they hold “with probability greater than $1 - p^{2-c}$ ”. For example, since $n > 2C_2^{-1}A_\Sigma^2 d_\delta^2 (c \log p + \log C_1)$, we have $A_\Sigma d_\delta \epsilon < 1/2$ with probability greater than $1 - p^{2-c}$ and we shall repeatedly use this inequality without mentioning it holds with probability greater than $1 - p^{2-c}$. Since $n > C_{2\delta}^{-1}(c \log p + \log C_1)$, by (17) of Ravikumar et al. (2011) and Theorem 1, we also have $\max\{\epsilon, \epsilon_\mu, \epsilon_\Omega\} \leq (C_3 + 1)\{(c \log p + \log C_1)/C_{2\delta}n\}^{1/2} := \epsilon_0$.

From Lemma 4 we have,

$$\begin{aligned}
&|\hat{\Sigma}_{e,D}\hat{\Sigma}_{D,D}^{-1} - \Sigma_{e,D}\Sigma_{D,D}^{-1}|_1 \\
&\leq |(\hat{\Sigma}_{e,D} - \Sigma_{e,D})\Sigma_{D,D}^{-1}|_1 + |\Sigma_{e,D}(\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1})|_1 + |(\hat{\Sigma}_{e,D} - \Sigma_{e,D})(\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1})|_1 \\
&\leq |\hat{\Sigma}_{e,D} - \Sigma_{e,D}|_1 |\Sigma_{D,D}^{-1}|_{1,\infty} + |\Sigma_{e,D}|_1 |\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1}|_{1,\infty} \\
&\quad + |\hat{\Sigma}_{e,D} - \Sigma_{e,D}|_1 |\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1}|_{1,\infty} \\
&\leq d_\delta \epsilon A_\Sigma + d_\delta (B + \epsilon) \|\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1}\|_{1,\infty} \\
&\leq d_\delta \epsilon A_\Sigma + \frac{d_\delta (B_\Sigma + d_\delta \epsilon) A_\Sigma^2 \epsilon}{1 - A_\Sigma d_\delta \epsilon}. \tag{22}
\end{aligned}$$

Combining (21), (22) and Lemma 5 we have:

$$\begin{aligned}
& |2\hat{\Sigma}_{e,D}\tilde{\delta}_D - \hat{\gamma}_e| \\
\leq & \left\{ \epsilon A_\Sigma + \frac{(B + d_\delta \epsilon) A_\Sigma^2 \epsilon}{1 - A_\Sigma d_\delta \epsilon} \right\} A_\gamma d_\delta + \left(2 - \alpha_\delta + d_\delta \epsilon A_\Sigma + \frac{d_\delta (B_\Sigma + d_\delta \epsilon) A_\Sigma^2 \epsilon}{1 - A_\Sigma d_\delta \epsilon} \right) K_\gamma \\
& + \lambda_\delta \left(1 - \alpha_\delta + d_\delta \epsilon A_\Sigma + \frac{d_\delta (B_\Sigma + d_\delta \epsilon) A_\Sigma^2 \epsilon}{1 - A_\Sigma d_\delta \epsilon} \right),
\end{aligned}$$

where

$$K_\gamma = 2\epsilon_0[4 + (2 + B_\Sigma A_2)|\Omega(\mu_1 - \mu_2)|_1 + 2B(A_1 + C_3)],$$

and we have used the fact that $\epsilon < B$, $d_\delta \epsilon A_2 < 1$, $d\epsilon < 1$ and hence $8\epsilon_\mu + 2(\epsilon + B_\Sigma \epsilon_\Omega A_2 + d\epsilon \epsilon_\Omega A_2)|\Omega(\mu_1 - \mu_2)|_1 + 2(B + \epsilon)(A_1 + d\epsilon_\Omega)\epsilon_\mu < K_\gamma$. Assume that

$$d_\delta \epsilon A_\Sigma + \frac{d_\delta (B_\Sigma + d_\delta \epsilon) A_\Sigma^2 \epsilon}{1 - A_\Sigma d_\delta \epsilon} \leq C_\delta \alpha_\delta \min\{\lambda, 1\}, \quad (23)$$

where $C_\delta = \frac{\alpha_\delta \lambda_\delta - (2 - \alpha_\delta) K_\gamma}{\alpha_\delta \lambda_\delta (1 + A_\gamma + K_\gamma)}$. It can be easily shown that $0 < C_\delta < 1$. We then have

$$\begin{aligned}
|2\hat{\Sigma}_{e,D}\tilde{\delta}_D - \hat{\gamma}_e| & \leq \lambda_\delta A_\gamma C_\delta \alpha_\delta + (2 - \alpha_\delta) K_\gamma + C_\delta \alpha_\delta \lambda K_\gamma + (1 - \alpha_\delta + C_\delta \alpha_\delta) \lambda \\
& = \lambda.
\end{aligned}$$

Next we complete the Proof of this part by showing that (23) holds.

Since $n > C_2^{-1} C_\delta^{-2} \alpha_\delta^{-2} d_\delta^2 (5A_\Sigma + 2B_\Sigma A_\Sigma^2)^2 \cdot (c \log p + \log C_1)$, we have

$$d_\delta \epsilon A_\Sigma + \frac{d_\delta (B_\Sigma + d_\delta \epsilon) A_\Sigma^2 \epsilon}{1 - A_\Sigma d_\delta \epsilon} \leq d_\delta \epsilon A_\Sigma + 4d_\delta A_\Sigma \epsilon + 2B_\Sigma A_\Sigma^2 d_\delta \epsilon \leq C_\delta \alpha_\delta.$$

On the other hand, since $\lambda \geq \frac{d_\delta (5A_\Sigma + 2B_\Sigma A_\Sigma^2)}{C_\delta \alpha_\delta} \epsilon_0$, we have

$$d_\delta \epsilon A_\Sigma + \frac{d_\delta (B_\Sigma + \epsilon d_\delta) A_\Sigma^2 \epsilon}{1 - A_\Sigma d_\delta \epsilon} \leq \lambda C_\delta \alpha_\delta.$$

(ii) Use the fact that $A_\Sigma d_\delta \epsilon < 1/2$ we have:

$$\begin{aligned}
|\hat{\delta}_D - \delta_D|_\infty &= \frac{1}{2} |\hat{\Sigma}_{D,D}^{-1}(\hat{\gamma}_D - \lambda_\delta Z_D) - \Sigma_{D,D}^{-1} \gamma_D| \\
&\leq |(\hat{\Sigma}_{D,D}^{-1} - \Sigma_{D,D}^{-1})\hat{\gamma}_D|_\infty + |\Sigma_{D,D}^{-1}(\hat{\gamma}_D - \gamma_D)|_\infty + \lambda_\delta |\hat{\Sigma}_{D,D}^{-1}|_{1,\infty} \\
&\leq \frac{A_\Sigma^2 d_\delta \epsilon}{1 - A_\Sigma d_\delta \epsilon} (A_\gamma + |\hat{\gamma}_D - \gamma_D|_\infty) + A_\Sigma |\hat{\gamma}_D - \gamma_D|_\infty + \lambda_\delta \left(A_\Sigma + \frac{A_\Sigma^2 d_\delta \epsilon}{1 - A_\Sigma d_\delta \epsilon} \right) \\
&\leq \frac{A_\gamma A_\Sigma^2 d_\delta \epsilon}{1 - A_\Sigma d_\delta \epsilon} + K_\gamma \left[\frac{A_\Sigma^2 d_\delta \epsilon}{1 - A_\Sigma d_\delta \epsilon} + A_\Sigma \right] + \lambda_\delta \left(A_\Sigma + \frac{A_\Sigma^2 d_\delta \epsilon}{1 - A_\Sigma d_\delta \epsilon} \right) \\
&\leq 2A_\gamma A_\Sigma^2 d_\delta \epsilon + 2K_\gamma A_\Sigma + 2\lambda_\delta A_\Sigma.
\end{aligned}$$

This theorem is proved by plugging in $K_\gamma = 2\epsilon_0[4 + (2 + B_\Sigma A_2)|\Omega(\mu_1 - \mu_2)|_1 + 2B(A_1 + C_3)]$.

Proofs of Theorem 3

Proof. (i) Write $D(z) = (z - \mu)^T \Omega (z - \mu) + \delta^T (z - \mu)$ and $\hat{D}(z) = (z - \hat{\mu})^T \hat{\Omega} (z - \hat{\mu}) + \hat{\delta}^T (z - \hat{\mu})$.

$$\begin{aligned}
R_n(1|2) &= P(\hat{D}(z) + \eta > 0 | z \sim N(\mu_2, \Sigma_2)) \\
&= P(D(z) + \eta > D(z) - \hat{D}(z) | z \sim N(\mu_2, \Sigma_2)).
\end{aligned}$$

Denote $z = (z_1, \dots, z_p)^T$. Note that $z^T(\Omega - \hat{\Omega})z \leq \sum_{ij} |z_i z_j| (\Omega - \hat{\Omega})_{ij}$, by noticing that $E|z_i z_j| \leq (Ez_i^2 + Ez_j^2)/2 \leq C_\mu^2 + C_\Sigma$, we have $z^T(\Omega - \hat{\Omega})z = O_p(s \|\Omega - \hat{\Omega}\|_\infty)$. Using a similar argument for bounding $|z^T \Omega \mu - z \hat{\Omega} \hat{\mu}|$, $|\mu^T \Omega \mu - \hat{\mu}^T \hat{\Omega} \hat{\mu}|$, $(\delta - \hat{\delta})^T z$ and $\delta^T \mu - \hat{\delta}^T \hat{\mu}$ we obtain,

$$|D(z) - \hat{D}(z)| = O_p \left(s C_3 \sqrt{\frac{c \log p + \log C_1}{C_2 n}} + d_\delta C_{3\delta} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}} \right). \quad (24)$$

From Assumption 1 and (24) and the mean value theorem, we have:

$$\begin{aligned}
R_n(1|2) - R(1|2) &= \int_0^{D(z) - \hat{D}(z)} F_2(z) dz \\
&= O_p \left(s C_3 \sqrt{\frac{c \log p + \log C_1}{C_2 n}} + d_\delta C_{3\delta} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}} \right).
\end{aligned}$$

(i) is proved by noticing that the above equality is also true for $R_n(2|1) - R(2|1)$.

(ii) Let $\Phi(\cdot)$ be the cumulative distribution function of a standard normal random variable. We have any constant $C_z > 0$,

$$P(|z|_\infty > C_z \sqrt{\log p}) \leq p \left[1 - \Phi \left(\frac{C \sqrt{\log p} + C_\mu}{C_\Sigma^{1/2}} \right) \right].$$

From Lemma 11 of Liu, Lafferty, and Wasserman (2009) we have when p is large enough, by choosing $C_z > \sqrt{2(c-1)}C_\Sigma^{1/2}$,

$$p \left[1 - \Phi \left(\frac{C_z \sqrt{\log p} + C_\mu}{C_\Sigma^{1/2}} \right) \right] \leq p^{2-c}.$$

This together with Theorems 1 and 2 and the proof in (i), we have with probability greater than $1 - 3p^{2-c}$,

$$|D(z) - \hat{D}(z)| = O_p \left(sC_3 \log p \sqrt{\frac{c \log p + \log C_1}{C_2 n}} + d_\delta C_{3\delta} \sqrt{\log p} \sqrt{\frac{c \log p + \log C_1}{C_{2\delta} n}} \right).$$

The rest of the proof is similar to that in the proof of (i). □

Proof of Proposition 2

Proof. Suppose $e_1 > e_2 \geq \eta$. Denote $c_i = |\Sigma_1 \Sigma_2^{-1}|^{1/2} \exp\{\frac{1}{2}[e_i - \frac{1}{4}(\mu_1 - \mu_2)^T \Omega(\mu_1 - \mu_2)]\}$ for $i = 1, 2$. We have,

$$\begin{aligned} R(d, e_i) &= \pi_1 \int_{D(z, e_1) < 0} f_1(z) dz + \pi_2 \int_{D(z, e_i) > 0} f_2(z) dz \\ &= \int_{f_1(z)/f_2(z) < c_i^{-1}} \pi_1 f_1(z) dz + \int_{f_1(z)/f_2(z) > c_i^{-1}} \pi_2 f_2(z) dz. \end{aligned}$$

Since $e_1 > e_2 \geq \eta$, it can be easily shown that $c_1 > c_2 \geq \pi_1/\pi_2$. Consequently we have

$$R(d, e_1) - R(d, e_2) = - \int_{c_1^{-1} < f_1(z)/f_2(z) < c_2^{-1}} [\pi_1 f_1(z) - \pi_2 f_2(z)] dz > 0.$$

Therefore, $R(d, e)$ is strictly monotone increasing on $e \in [\eta, \infty)$. The second statement can be similarly proved. $\square$

Proofs of Theorem 4

We first introduce some technical lemmas and the the proof of Theorem 4 will be given after these lemmas.

For any constant c , define $l_c = \min\{\text{ess inf}_{z \in (-c, c)} F_i(z), i = 1, 2\}$.

Lemma 6. *For any $-c < \epsilon_\eta < c$, we have $R(d, \eta + \epsilon_\eta) - R(d, \eta) \leq \pi_2 u_c |\epsilon_\eta|$ and $R(d, \eta + \epsilon_\eta) - R(d, \eta) \geq \epsilon_\eta^2 \exp(-c/2) \pi_2 l_c / 4$.*

Proof. Let's consider $0 < \epsilon_\eta \leq c$ first. Note that

$$R(d, \eta + \epsilon_\eta) - R(d, \eta) = \int_{-\epsilon_\eta < D(z, \eta) < 0} [\pi_2 f_2(z) - \pi_1 f_1(z)] dz.$$

We have

$$R(d, \eta + \epsilon_\eta) - R(d, \eta) \leq \int_{-\epsilon_\eta < D(z, \eta) < 0} \pi_2 f_2(z) dz = \pi_2 \int_{-\epsilon_\eta}^0 F_2(z) dz \leq \pi_2 u_c \epsilon_\eta.$$

By noticing that $1 - \exp(-x/2) - x \exp(-c/2)/2$ is an increasing function in $[0, c]$ we have

$$\begin{aligned} R(d, \eta + \epsilon_\eta) - R(d, \eta) &> \int_{-\epsilon_\eta < D(z, \eta) < -\epsilon_\eta/2} [\pi_2 f_2(z) - \pi_1 f_1(z)] dz \\ &> \int_{-\epsilon_\eta < D(z, \eta) < -\epsilon_\eta/2} [1 - \exp(-\epsilon_\eta/2)] \pi_2 f_2(z) dz \\ &> \int_{-\epsilon_\eta < D(z, \eta) < -\epsilon_\eta/2} \epsilon_\eta \exp(-c/2) \pi_2 f_2(z) / 2 dz \\ &> \epsilon_\eta^2 \exp(-c/2) \pi_2 l_c / 4. \end{aligned}$$

The lemma is then proved using a same argument as above for $-c < \epsilon_\eta \leq 0$. $\square$

Similar to the proof of Lemma 6 where the bounds do not depend on η , we have:

Lemma 7. *Suppose $\hat{d} - d = O_p(\Delta_d)$ with $\Delta_d \rightarrow 0$, we have $R(\hat{d}, e) - R(d, e) = O_p(\Delta_d u_c)$ uniformly in $e \in (-c, c)$.*

Lemma 8. *Under the assumptions of Theorem 3, $R_n(\hat{d}, e) \rightarrow R(d, e)$ in probability uniformly in $e \in (-c, c)$.*

Proof. Denote all the samples in the two classes as $\{z_i, i = 1, \dots, n_1 + n_2\}$ and denote the estimator obtained by leaving the i th sample out as $\hat{d}_{-i}$. Similarly we use $\hat{d}_{-(i,j)}$ to denote the estimator obtained by leaving the i th and j th sample out. From (24), we immediately have that for any $e \in (-c, c)$,

$$\begin{aligned} & EI\{\hat{d}(z_i) + e > 0\} - EI\{\hat{d}_{-i}(z_i) + e > 0\} \\ &= EI\{d(z_i) + e > 0\} - EI\{d(z_i) + e > 0\} + o(1) \\ &= o(1). \end{aligned}$$

Together with Lemma 7 we have

$$ER_n(\hat{d}, e) - R(d, e) = ER(\hat{d}_{-i}, e) + o(1) - R(d, e) \rightarrow 0, \quad (25)$$

uniformly in $e \in (-c, c)$. Note that

$$\text{Var}(I\{\hat{d}(z_i) + e > 0\}) \leq \frac{1}{4},$$

and for any $(i, j) \in \{(k, l) : 1 \leq k, l \leq n_1 + n_2, i \neq j\}$,

$$\begin{aligned}
& Cov(I\{\hat{d}(z_i) + e > 0\}, I\{\hat{d}(z_j) + e > 0\}) \\
&= Cov(I\{\hat{d}_{-(i,j)}(z_i) + e > 0\}, I\{\hat{d}_{-(i,j)}(z_j) + e > 0\}) + o(1) \\
&= Cov(I\{d(z_i) + e > 0\}, I\{d(z_j) + e > 0\}) + o(1),
\end{aligned}$$

where the last step can be obtained using (24) and Lemma 7 and the $o(1)$ term does not depend on e . Since z_i, z_j are independent, we immediately have

$$Var(R_n(\hat{d}, e)) \rightarrow 0, \tag{26}$$

uniformly in $e \in (-c, c)$. The lemma is then proved by Markov's inequality and the uniform convergence of the bias (25) and the variance (26) of $R_n(\hat{d}, e)$. $\square$

Proof of Theorem 4

The result that $\hat{\eta} \rightarrow \eta$ can be obtained by Proposition 2, Lemma 8 and Theorem 5.7 of Van der Vaart (2011). The second statement immediately follows from Theorem 3.